

=> fil wpix

FILE 'WPIX' ENTERED AT 11:48:13 ON 22 JUL 2003  
COPYRIGHT (C) 2003 THOMSON DERWENT

FILE LAST UPDATED: 19 JUL 2003 <20030719/UP>  
MOST RECENT DERWENT UPDATE: 200346 <200346/DW>  
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> NEW WEEKLY SDI FREQUENCY AVAILABLE --> see NEWS <<<

>>> SLART (Simultaneous Left and Right Truncation) is now  
available in the /ABEX field. An additional search field  
/BIX is also provided which comprises both /BI and /ABEX <<<

>>> PATENT IMAGES AVAILABLE FOR PRINT AND DISPLAY <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES,  
SEE <http://www.derwent.com/dwpi/updates/dwpicov/index.html> <<<

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,  
PLEASE VISIT:  
[http://www.stn-international.de/training\\_center/patents/stn\\_guide.pdf](http://www.stn-international.de/training_center/patents/stn_guide.pdf) <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER  
GUIDES, PLEASE VISIT:  
[http://www.derwent.com/userguides/dwpi\\_guide.html](http://www.derwent.com/userguides/dwpi_guide.html) <<<

=> d all abeq tech abex tot l116

L116 ANSWER 1 OF 5 WPIX COPYRIGHT 2003 THOMSON DERWENT on STN

AN 2003-148342 [14] WPIX

DNC C2003-038281

TI Synthesis of proteins using cell-free protein system, e.g. membrane  
proteins, receptors, or channel proteins, useful for developing drugs,  
comprises adding surfactant to the system to produce the protein without  
agglutination.

DC B04 D16

IN ISHIHARA, G; KIGAWA, T; SHIROUZU, M; TAJIMA, K; YABUKI, T;  
YOKOYAMA, S

PA (RIKE) RIKEN KK

CYC 22

PI WO 2002090537 A1 20021114 (200314)\* JA 45p C12N015-09 <--  
RW: AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE TR  
W: CA US

JP 2003018999 A 20030121 (200317) 15p C12P021-02 <--

ADT WO 2002090537 A1 WO 2002-JP4204 20020426; JP 2003018999 A JP 2001-285145  
20010919

PRAI JP 2001-285145 20010919; JP 2001-135111 20010502

IC ICM C12P021-02

ICS C07K001-02; C12N001-21

ICA C12N015-09

AB WO 2002090537 A UPAB: 20030227

NOVELTY - Synthesis (M1) of a protein by using a cell-free protein  
synthesis system comprises adding a surfactant to the system so that the  
protein can be produced without agglutination.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the  
following:

(1) A similar method (M2) in which an extractant for bacterial mass  
is applied as well;

(2) Restructuring (M3) a protein produced in a cell-free protein  
synthesis system containing a template nucleic acid encoding at least a  
part of a membrane protein, surfactant and lipid, in which and  
concentration of the surfactant in the synthetic reaction solution is

lowered during the protein synthesis or a definite period of time afterwards; and

(3) Synthesizing (M4) a membrane protein that has at least a part of its biological activity comprising:

(a) synthesis of the protein in a cell-free protein synthesis system containing a cell extraction solution, a template nucleic acid encoding at least a part of a membrane protein, surfactant and lipid; and

(b) lowering concentration of the surfactant in the synthetic reaction solution during the protein synthesis or after a definite period of time.

USE - The method is useful for synthesis of proteins, restructuring proteins, and synthesizing a membrane protein (claimed), especially proteins with hydrophobic domains e.g. membrane proteins, receptors, channel proteins or transporters, which are then applicable in studying their structure and function and in developing drugs to control such functions and in diagnosis and treatment of related diseases.

ADVANTAGE - Such method is efficient and quantitative, without loss of protein activity due to problems in agglutination of difficult to dissolve proteins and in the folding process, on a large scale.

Dwg.0/11

FS CPI

FA AB; DCN

MC CPI: B01-A03; B04-C03C; B04-E01; B04-E03F; B04-K0100E; B04-L0100E; B04-N0400E; B05-B01P; B07-A02B; B10-A22; B10-D03; B10-E04C; B11-C08; B12-K04A; B12-K04E; D05-H09; D05-H12A; D05-H17

TECH UPTX: 20030227

TECHNOLOGY FOCUS - BIOTECHNOLOGY - Preferred Processes: The protein is a protein containing at least a hydrophobic domain, which is particularly a membrane protein, including receptor, channel protein, transporter and membrane-binding enzyme, or their fragment.

TECHNOLOGY FOCUS - ORGANIC CHEMISTRY - Preferred Processes: Such surfactant is a mild surfactant which will not denature the protein, e.g. a non-ionic or amphoteric surfactant including digonin, polyoxyethylene alkyl ether, polyoxyethylene sorbitan, beta-dodecyl maltoside, beta-octyl glucoside, beta-nonyl glucoside, beta-heptyl thioglucoside, beta-octyl thioglucoside, sucrose monodecanoate, sucrose monodecanoate, octyltetraoxyethylene, octylpentaoxyethylene, dodecylloctaoxyethylene, N,N-dimethyldodecylamine N-oxide, N,N-dimethyldodecylamine N-oxide, N,N-dimethyldodecylammoniopropanesulfonate, octyl(hydroxyethyl) sulfoxide, octanoyl-N-methylglucamide, nonanoyl-N-methylglucamide, decanoyl-N-methylglucamide and 3-((3-cholamidopropyl)-dimethylammonio)-1-propanesulfonate (CHAPS). Concentration of such surfactant is particularly 0.1-2 volume % digonin and/or 0.01-0.5 volume % Brij 35. To lower the surfactant concentration, dialysis, dilution, filtration, centrifugation and/or addition of an adsorbent for adsorption of such surfactant is applied.

ABEX UPTX: 20030227

EXAMPLE - To synthesize a rat neurotensin receptor (NRT), an MBP-T43NTR-TrxA-H10 template fragment was prepared after two PCR reactions, with Escherichia coli S30 extract (Zubay et al., Annu. Rev. Geneti., 1973, vol. 7, p. 267) for reaction in a cell-free protein synthesis system with e.g. 0.4% digonin added. The required protein was easily recovered.

L116 ANSWER 2 OF 5 WPIX COPYRIGHT 2003 THOMSON DERWENT on STN

AN 2003-140656 [13] WPIX

DNN N2003-111659 DNC C2003-035853

TI Structural analysis of proteins useful for identifying proteins, their domain structures and global stereostructures, comprises synthesis of protein with tag sequence and measuring by mass spectrometry.

DC B04 D16 S03

IN KIGAWA, T; SHEN, X; YOKOYAMA, S

PA (RIKA) RIKAGAKU KENKYUSHO; (RIKE) RIKEN KK

CYC 22

PI WO 2002099437 A1 20021212 (200313)\* JA 34p G01N033-68 <--

RW: AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE TR

W: CA US

JP 2002372517 A 20021226 (200314) 13p G01N027-62 <--

ADT WO 2002099437 A1 WO 2002-JP4473 20020508; JP 2002372517 A JP 2001-164819  
20010531

PRAI JP 2001-164819 20010531

IC ICM G01N027-62; G01N033-68

ICS C12N015-09; C12Q001-37; G01N033-483;

G01N033-53

AB WO 200299437 A UPAB: 20030224

NOVELTY - Identifying the molecular weight of a protein by mass spectrometry comprises:

(a) synthesizing a protein fused with a tag sequence, in a cell-free protein synthesis system;

(b) contacting the fused protein with a tag sequence-binding carrier;

(c) collecting the bonded carrier; and

(d) measuring molecular weight of the fused protein thus collected.

DETAILED DESCRIPTION - Identifying (M1) the molecular weight of a protein by mass spectrometry (MS) comprises:

(a) synthesizing a protein fused with a tag sequence in a cell-free protein synthesis system;

(b) contacting the fused protein with a carrier capable of binding to the tag sequence;

(c) collecting the carrier bonded to the fused protein; and

(d) measuring molecular weight of the fused protein by MS with use of the fused protein bonded to the carrier as collected above.

INDEPENDENT CLAIMS are also included for the following:

(1) Identifying (M2) a protein domain by MS comprising:

(a) synthesizing a protein fused with a tag sequence in a cell-free protein synthesis system;

(b) contacting the fused protein with a carrier capable of binding to the tag sequence;

(c) collecting the carrier bonded to the fused protein;

(d) partial decomposition of the fused protein bonded to the carrier as collected above by protease; and

(e) measuring molecular weight of the fused protein and/or decomposition products by MS with use of solution of the decomposition reaction; and

(2) Identifying (M3) a protein domain by MS comprising:

(a) synthesizing a protein fused with a tag sequence in a cell-free protein synthesis system;

(b) contacting the fused protein with a carrier capable of binding to the tag sequence;

(c) collecting the carrier bonded to the fused protein;

(d) measuring molecular weight of the fused protein by MS with use of the fused protein bonded to the carrier as collected above;

(e) partial decomposition of the fused protein bonded to the carrier as collected by protease, and measuring molecular weight of the decomposition product by MS with use of solution of the decomposition reaction; and

(f) comparing such measured molecular weight of the fused protein and that of the decomposition product.

USE - The analytical method is applicable in identifying the molecular weight of proteins (claimed), their domain structures and studying other global stereostructures, and structural genomic investigation.

ADVANTAGE - With such method, synthesis, purification and analysis of a protein can be efficiently and quickly carried out, particularly on multi-specimen screening plates consecutively with automation, in a high-throughput manner as stably adsorbed entity, without problems due to

the insoluble purified protein.

Dwg.0/4

FS CPI EPI

FA AB; DCN

MC CPI: B04-B03C; B04-C01; B04-G01; B04-K01; B04-N0400E; B11-C08A; B11-C08D;  
B11-C08E3; B11-C08F2; B12-K04E; D05-A02C; D05-H09; D05-H11; D05-H17  
EPI: S03-E10; S03-E14H; S03-E14H4

TECH UPTX: 20030224

TECHNOLOGY FOCUS - ORGANIC CHEMISTRY - Preferred Method: The carrier capable of binding specifically to the tag sequence is a carrier holding 1 or more selected from antibodies, binding proteins, receptors and parts containing their binding regions; or chosen from metal atoms, amino acids, peptides, oligonucleotides and heterocyclic compounds. The MS can be time-of-flight type matrix-assisted laser-desorption ionization technique (MALDI-TOF).

ABEX UPTX: 20030224

EXAMPLE - Identifying the molecular weight of a protein by mass spectrometry was carried out as follows. A His-Ras protein was synthesized in a cell-free protein synthesis system (Annu. Rev. Genet., 1973, vol. 7, p. 267, then purified on nickel-bound agarose-beads and MALDI-TOF MS analysis of the agarose bead-adsorbed Ras protein, with observation of (Ras)+ peak at m/z = 22108.1 and (Ras)2+ peak at m/z 11053.5.

L116 ANSWER 3 OF 5 WPIX COPYRIGHT 2003 THOMSON DERWENT on STN

AN 2003-138616 [13] WPIX

DNC C2003-035267

TI Producing a soluble protein domain comprises expressing a fusion protein that is coded on each of the DNA fragments fused with a DNA encoding a functional protein and synthesizing the soluble protein domain in a **cell-free** system.

DC B04 D16

IN KIGAWA, T; SEKI, E; YOKOYAMA, S

PA (RIKE) RIKEN KK; (KIGA-I) KIGAWA T; (SEKI-I) SEKI E; (YOKO-I) YOKOYAMA S

CYC 3

PI US 2002142387 A1 20021003 (200313)\* 12p C12P021-02 <--  
CA 2362784 A1 20020906 (200313) EN C12N015-11 <--  
JP 2002262873 A 20020917 (200313) 9p C12N015-09 <--

ADT US 2002142387 A1 US 2001-994573 20011126; CA 2362784 A1 CA 2001-2362784  
20011123; JP 2002262873 A JP 2001-62703 20010306

PRAI JP 2001-62703 20010306

IC ICM C12N015-09; C12N015-11; C12P021-02

ICS C07K014-00; C07K014-47; C07K019-00;  
C12N001-21; C12N015-62; C12N015-70;  
C12P019-34; C12P021-00; C12Q001-68;  
G01N023-20; G01N024-08; G01N033-566;  
G01N033-68

ICI C12P021-02; C12R001:19

AB US2002142387 A UPAB: 20030224

NOVELTY - Producing a soluble protein domain comprising preparing two or more DNA fragments by partially digesting a DNA coding for a protein, expressing a fusion protein that is coded on each of the DNA fragments fused with a DNA encoding a functional protein exhibiting a function, selecting the fusion protein exhibiting the function, and synthesizing the soluble protein domain in a **cell-free** system, is new.

DETAILED DESCRIPTION - An INDEPENDENT CLAIM is also included for a method for analyzing the three dimensional structure of a protein.

USE - The method is useful for producing a soluble protein domain (claimed).

Dwg.0/5

FS CPI

FA AB; DCN

MC CPI: B04-N0400E; B04-N08; B11-C08G1; B11-C08G2; B12-K04E; D05-H09;  
D05-H17C

TECH

UPTX: 20030224

TECHNOLOGY FOCUS - BIOTECHNOLOGY - Preferred Method: Producing a soluble protein domain further comprises:

- (a) selecting a fusion protein from the fusion proteins containing a functional protein exhibiting a function; and
- (b) synthesizing a soluble protein domain from the fusion protein selected from step (a).

The method also comprises:

- (a) constructing an expression vector that expresses a fusion protein with a **green fluorescent** protein or its derivative, where the expression vector comprises a DNA coding for a protein and a gene for the **green fluorescent** protein or its derivative;
- (b) preparing two or more DNA fragments by partially digesting the expression vector with DNA decomposition enzyme;
- (c) transforming Escherichia coli with each of the DNA fragments to obtain two or more transformed E. coli isolating a transformed clone that emits fluorescence among the transformed E. coli;
- (d) recovering the DNA from the isolated transformed clone; and
- (e) synthesizing the soluble protein domain that is coded on the recovered DNA in a **cell-free** system.

The method may also comprise:

- (a) preparing an expression vector comprising a DNA coding for a protein and a DNA coding for a functional protein;
- (b) treating the vector by using a decomposing enzyme and forming two or more vectors, each vector comprising a fragment of the DNA coding for a protein and the DNA coding for a functional protein;
- (c) expressing a fusion protein that is coded on each of the vectors fused with a DNA encoding a functional protein;
- (d) selecting the fusion protein exhibiting the function among two or more fusion proteins synthesized in step (a);
- (e) synthesizing the soluble protein domain in a **cell-free** system.

The functional protein consists of enzyme, binding protein, luminescent protein, fluorescent protein or its functional protein. The fluorescent protein is a **green fluorescent** protein or its derivative. The selection is performed by:

- (a) transforming a recipient **cell** with each of the DNA fragments and the DNA of the functional protein; and
- (b) selecting the clone that exhibits the function in the obtained transformants.

The selection may also be performed by measuring the function of the expressed proteins. The recipient **cell** is E. coli. Analyzing the three dimensional structure of a protein comprises:

- (a) synthesizing the soluble protein domain by the method of producing a soluble protein domain; and
- (b) analyzing the three dimensional structure of the soluble protein domain by X-ray crystallography or NMR spectroscopy.

L116 ANSWER 4 OF 5 WPIX COPYRIGHT 2003 THOMSON DERWENT on STN

AN 2001-257972 [26] WPIX

DNC C2001-077772

TI New isolated, enriched, **cell free** and/or recombinant nucleic acid useful for e.g. altering **cell** proliferation characteristic such as to alter plant **cell**, organ or tissue size.

DC C06 D16

IN GUTIERREZ-ARMENTA, C; RAMIREZ-PARRA, E

PA (CNSJ) CONSEJO SUPERIOR INVESTIGACIONES CIENTIF

CYC 93

PI WO 2001021644 A2 20010329 (200126)\* EN 77p C07K014-00 <--

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ

NL OA PT SD SE SL SZ TZ UG ZW

W: AE AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CR CU CZ DE DK DM EE

ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR  
 LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD SE SG SI  
 SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

AU 2000077814 A 20010424 (200141) C07K014-00 <--  
 EP 1214436 A2 20020619 (200240) EN C12N015-82 <--  
 R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT  
 RO SE SI

BR 2000014634 A 20030225 (200320) C07K014-00 <--  
 JP 2003510040 W 20030318 (200321) 89p C12N015-09 <--  
 ES 2182615 A1 20030301 (200327) C07K014-415 <--

ADT WO 2001021644 A2 WO 2000-EP9325 20000925; AU 2000077814 A AU 2000-77814  
 20000925; EP 1214436 A2 EP 2000-967766 20000925, WO 2000-EP9325 20000925;  
 BR 2000014634 A BR 2000-14634 20000925, WO 2000-EP9325 20000925; JP  
 2003510040 W WO 2000-EP9325 20000925, JP 2001-525218 20000925; ES 2182615  
 A1 ES 1999-2127 19990924

FDT AU 2000077814 A Based on WO 200121644; EP 1214436 A2 Based on WO  
 200121644; BR 2000014634 A Based on WO 200121644; JP 2003510040 W Based on  
 WO 200121644

PRAI ES 1999-2474 19991111; ES 1999-2127 19990924

IC ICM C07K014-00; C07K014-415; C12N015-09;  
 C12N015-82

ICS A01H005-00; C07K016-16; C12N005-04; C12N005-10; C12N005-14;  
 C12N015-11; C12N015-29; C12N015-62;  
 C12Q001-68

AB WO 200121644 A UPAB: 20010515

NOVELTY - An isolated, enriched, **cell free** and/or  
 recombinant nucleic acid (II) comprising a sequence encoding for  
 expression of a protein or peptide (I) capable of altering  
 E2F-dimerization partner (DP) activity in a plant **cell** is new.

DETAILED DESCRIPTION - (P1) is characterized in that it has one or  
 both DP activities in plants selected from:

- (1) the ability to dimerized with plant E2F protein; and
- (2) the ability to modulate, particularly enhance, E2F binding to E2F  
 transcription factor binding sites in plant DNA or effect of it  
 characterized in that the protein or peptide comprises a 261 amino acid  
 sequence (S1) or a functionally active part of it or a sequence having at  
 least 70% homology to so the sequence or part of it.

INDEPENDENT CLAIMS are also included for the following:

- (1) controlling one or more of plant growth, gene expression,  
 cellular DNA replication, cell cycle progression, differentiation and  
 development comprising increasing or decreasing DP protein activity in a  
 plant cell through the expression of (P1);
- (2) isolated (I);
- (3) a nucleic acid probe (III) comprising a DNA sequence  
 corresponding to an amino acid sequence comprising 261, 458, 52, or 51  
 amino acids fully defined in the specification;
- (4) a nucleic acid probe or primer (IV) comprising a double or single  
 stranded DNA of sequence corresponding to 10 or more contiguous  
 nucleotides taken from a sequence (S2) comprising 1089 base pairs fully  
 defined in the specification, with the proviso that it is not selected  
 from residues 70-136;
- (5) an oligonucleotides probe (V) comprising at least 18 contiguous  
 bases of S2;
- (6) an antisense DNA (VI) to (II), (III), (IV) or (V);
- (7) a nucleic acid vector or construct (VII) comprising (II), (III),  
 (IV), (V), or (VI) or its antisense nucleic acid;
- (8) a plant cell (IX) comprising (VIII);
- (9) a transgenic plant or its part (X) comprising (IX);
- (10) an antibody (XI) raised against (I);
- (11) identifying and/or isolating (M2) DNAs corresponding to complete  
 or partial genes that are regulated in G1 passage, G1/S phase transition  
 and/or S phase progression of the cell cycle, involves contacting sample  
 of genomic DNA with binding material specific for binding such complete or

partial genes, removing non-bound DNA from the specific binding material, and releasing and isolating the bound DNA, where the specific binding material comprises a peptide or protein including the DNA binding sequence of a protein that is capable of acting as a part of a plant hetero-oligomer transcription activator or repressor; and

(12) a specific binding material (XII) which comprises a peptide or protein having DNA binding activity with respect to plant DNA transcription activator or repressor factor binding sites, particularly in genomic DNA, and having the ability to dimerize or oligomerize with a further such plant protein together with one or more of a further peptide or protein.

USE - (IV) is useful in Northern and Southern blotting and in polymerase chain reaction (PCR), including RT-PCR and LCR (claimed). (M1) or (II) is useful for altering plant cell, organ or tissue shape, and particularly to alter cell proliferation characteristic such as to alter plant cell, organ or tissue size. (M1) is useful for altering expression of other proteins with transformed cells and cells derived from it, particularly direct or indirect progeny. (II) is useful for obtaining sequences of (II) and, by using computational homology analysis of other organism's genome e.g. of plant, animal, yeast, bacteria, virus or fungal genomes, or molecular biology probing or other analysis techniques to identify therapeutic or otherwise industrially useful target genes involved in cell cycle that are not readily obtainable due to the comparative complexity or lack of information of/on such systems. (XI) is useful for identification and isolation of (I) and (II).

Dwg.0/3

FS CPI

FA AB; DCN

MC CPI: C04-A0800E; C04-C01G; C04-E03F; C04-E05; C04-E06; C04-E08;  
C04-F0800E; C04-G0100E; C04-N04A0E; C11-C07A; C11-C08E5; C12-K04E;  
C12-K04F; D05-H09; D05-H11; D05-H12A; D05-H12D1; D05-H12D2; D05-H12E;  
D05-H14B3; D05-H16B

TECH UPTX: 20010515

TECHNOLOGY FOCUS - BIOTECHNOLOGY - Preferred Method: The plant DP activity comprises one or both of the ability to dimerize with plant E2F protein, and the ability to modulate E2F binding to E2F/DP transcription factor binding sites in plant DNA. M1 comprises altering the plant DP protein level, the E2F-DP DNA binding activity, transactivation properties and/or the DP/E2F binding activity. DP may be modified alone and/or in combination with a modification of the levels or activity of plant E2F and/or plant Rb. In M2, the specific binding material comprises a peptide or protein which includes a plant E2F DNA binding domain, together with a plant E2F-DP hetero-dimerization domain and/or a plant retinoblastoma protein E2F binding domain. The E2F and DP domains are comprised in *Triticum monococcum* sequences or its functional variants or parts, or the DP domains are comprised in (II). The specific binding material peptide or protein is labeled or tagged to assist in identifying or immobilizing it, particularly when in bound complex with the gene to be identified and/or isolated, and comprises two peptides or proteins, one including the E2F DNA binding domain and one including the DP hetero-dimerization domain bound together as a hetero-dimer. The E2F sequence is all or part of a sequence comprising 458 amino acids fully defined in the specification or a sequence having 90% homology to it. Preferred Polypeptide: (I) has at least 50% or more, preferably 70% or more identity with S1, or its functional part. (I) of sequence S1, or its variant modified such that the amino acid sequence is mutated so that its ability to dimerize with E2F protein is reduced from that of the native sequence or abolished completely, where the peptide is capable of acting as a DP protein which decreases or abolishes native or recombinant E2F binding to its DNA binding site, thus inhibiting or abolishing E2F activity in a cell in which it is present. (I) has at least 70% homology to a sequence selected from: ARAAMAPPRGGAAAAATAALDLTG VHILEASSVPPLPERGGNAVQRKGAVDP, DKDRKKEKAAAPRITGWGLREYSKIVCEKVEAKGRTTYNEVADEIYSELKS,

MAHIGQGFDEKNIRRRVYDAFNVLIALRVIACEKKEIRWMGLSNRYEYKIKKLEEV, RKELVNKIRNKKALLQEI EKQFDDLQNIKLNRQTLESSAENVNGIRLPFVLVKT SR, and KARVEIEISDDSKFAHFEFNGAPFTLHDDL SILEGVRGNSIGKAGRATLH, fully defined in the specification. Preferred Polynucleotide: (II) has a sequence complementary or antisense to S2. (II) is contained in plasmid pCLON33, deposit number CECT 5195 made on August 17th 1999 under the terms of the Budapest Treaty for the International Recognition of Microorganism Deposits for Patent Purposes of 28th April 1977 at the Coleccion Espanola de Cultivos Tipo. (II) has a sequence encoding an E2F protein or peptide, and encodes (I) fused to a sequence encoding for a protein label. (II) is under the control of the same regulatory element or elements. (V) is 30-100, preferably 10-20 bases long. Preferred Antibody: (XI) is capable of specifically binding with plant DP factor peptides or proteins, but not to the human, mouse or Xenopus DP. Preferred Material: (XII) comprises a peptide or protein having DNA binding activity with respect to plant DNA E2F transcription factor binding sites, where the peptides or proteins are wheat peptides or proteins and its functional variants, or parts. (XII) is in the form of a coating or otherwise bound form on a support material, or in the form of particles of the peptides or proteins and/or hetero-dimers or trimers.

ABEX UPTX: 20010515

SPECIFIC SEQUENCES - (II) has a sequence S2 comprising 1089 base pairs fully defined in the specification (claimed).

EXAMPLE - In order to isolate protein partners for wheat (Triticum monococcum) TmE2F, E2F expressed from cDNA described in PCT/EP99/03158 was fused to the Gal4-DNA binding domain and applied to yeast two-hybrid screening. The TmE2F clone by itself transactivated the reporter genes and could not be used as bait. Therefore the last 85 amino acids of TmE2F (which, based on amino acid homology studies and binding experiments, should contain the Rb-binding and trans-activation domains but not a putative dimerization domain), were deleted. This C-terminally truncated TmE2F(1-373) did not trans-activate the reporter genes in yeast and was used as a bait in a screening of a wheat cDNA library constructed as a fusion to the Gal4-activation domain. The positive clones were allowed to grow, and thirty clones were isolated. Six clones with identical DNA sequence were identified and isolated by partial DNA sequencing. These sequences were used as a query in a BLAST search against GENBANK database, and the already identified members of the animal E2F-dimerization partner (DP) family were retrieved. The TmDP cDNA clone isolated by two-hybrid screen was 1089 base pairs long, including the poly-A tract, and contained a 5'-untranslated region and a 233 base pair 3'-untranslated region. The cDNA clone contained a single open reading frame of 261 amino acids.

L116 ANSWER 5 OF 5 WPIX COPYRIGHT 2003 THOMSON DERWENT on STN

AN 1991-132863 [18] WPIX

DNC C1991-057339

TI Cell free synthesis and isolation of novel genes - which are semi-random sequence and bind to e.g. antibodies or toxins.

DC B04 D16

IN KAWASAKI, G; KAWASAKI, G H

PA (KAWA-I) KAWASAKI G; (OPTE-N) OPTEIN INC

CYC 19

PI WO 9105058 A 19910418 (199118)\* 52p

RW: AT BE CH DE DK ES FR GB IT LU NL SE

W: AU CA JP KR NO

AU 9065373 A 19910428 (199131)

EP 494955 A1 19920722 (199230) EN 52p C12P021-00

R: AT BE CH DE DK ES FR GB IT LI LU NL SE

JP 05503000 W 19930527 (199326) 18p C12P021-00

AU 638762 B 19930708 (199334) C12N015-10

EP 494955 A4 19920812 (199523)

US 5643768 A 19970701 (199732) 14p C12P019-34

US 5658754 A 19970819 (199739) 12p C12P021-02



EP 494955 B1 19980715 (199832) EN C12P021-00  
R: AT BE CH DE DK ES FR GB IT LI LU NL SE  
DE 69032483 E 19980820 (199839) C12P021-00  
ES 2118066 T3 19980916 (199848) C12P021-00  
KR 204359 B1 19990615 (200063) C12N015-00  
KR 204360 B1 19990615 (200063) C12N015-00  
JP 3127158 B2 20010122 (200112) 17p C12P021-00  
KR 185192 B1 19990401 (200113) C12N015-10  
JP 2001078787 A 20010327 (200122) 17p C12N015-09  
CA 2067194 C 20030318 (200325) EN C12N015-10

ADT EP 494955 A1 EP 1990-915566 19901004, WO 1990-US5682 19901004; JP 05503000  
W JP 1990-514372 19901004, WO 1990-US5682 19901004; AU 638762 B AU  
1990-65373 19901004; EP 494955 A4 EP 1990-915566 ; US 5643768 A  
Cont of US 1989-417357 19891005, Div ex US 1991-798985 19911129, US  
1995-469026 19950606; US 5658754 A Cont of US 1989-417357 19891005, US  
1991-798985 19911129; EP 494955 B1 EP 1990-915566 19901004, WO 1990-US5682  
19901004; DE 69032483 E DE 1990-632483 19901004, EP 1990-915566 19901004,  
WO 1990-US5682 19901004; ES 2118066 T3 EP 1990-915566 19901004; KR 204359  
B1 WO 1990-US5682 19901004, Div ex KR 1992-700769 19920404, KR 1998-708672  
19981028; KR 204360 B1 WO 1990-US5682 19901004, Div ex KR 1992-700769  
19920404, KR 1998-708673 19981028; JP 3127158 B2 JP 1990-514372 19901004,  
WO 1990-US5682 19901004; KR 185192 B1 WO 1990-US5682 19901004, KR  
1992-700769 19920404; JP 2001078787 A Div ex JP 1990-514372 19901004, JP  
2000-240298 19901004; CA 2067194 C CA 1990-2067194 19901004, WO  
1990-US5682 19901004

FDT EP 494955 A1 Based on WO 9105058; JP 05503000 W Based on WO 9105058; AU  
638762 B Previous Publ. AU 9065373, Based on WO 9105058; EP 494955 B1  
Based on WO 9105058; DE 69032483 E Based on EP 494955, Based on WO  
9105058; ES 2118066 T3 Based on EP 494955; JP 3127158 B2 Previous Publ. JP  
05503000, Based on WO 9105058; CA 2067194 C Based on WO 9105058

PRAI US 1989-417357 19891005; US 1991-798985 19911129; US 1995-469026  
19950606

REP 2.Jnl.Ref; GB 2183661; No-Citns.; UK 2183661

IC C07H015-12; C07K007-00; C12N015-10; C12P019-34; C12P021-02  
ICM C12N015-00; C12N015-09; C12P019-34; C12P021-00; C12P021-02  
ICS C07H015-12; C07K007-00; C12N015-11; C12R001:19; C12R001:91;  
G01N033-15; G01N033-50; G01N033-566

ICA C12N015-10

ICI C12P021-00, C12R001:91

AB WO 9105058 A UPAB: 19940223  
Novel polypeptides (I) prodn. comprises construction of an in vitro  
expression unit (II) (contg. a 5' untranslated region with an RNA  
polymerase binding sequence, a ribosome binding sequence and a translation  
initiation region), attaching it to at least one semi-random nucleotide  
sequence and expressing or replication associated sequences followed by  
translation of the RNA produced to produce polysomes. The polysomes are  
bound to a substance of interest, isolated, disrupted to release the RNA  
which is recovered and converted to a cDNA which can be expressed as a  
novel polypeptide.

Also claimed are: 1. (I) where the RNA is translated into  
biologically active peptides or sub divided in such a way that the gene of  
interest is isolated and used to construct a cDNA to express the novel  
polypeptides; 2. (I) as above where the RNA is subdivided after  
translation into active polypeptides; 3. Isolation of a nucleotide  
sequence encoding a polypeptide of interest comprising transcribing (II)  
and at least one semi-random nucleotide to produce a mRNA library which is  
translated to produce polypeptides still attached to polysomes, and the  
polysomes are contacted to a substance of interest and the mRNA is  
recovered from the bound polysome.

USE - The method allows the isolation of semi-random DNA or RNA  
sequences that encode novel polypeptides which can bind specific  
substances. @ (52pp Dwg.No.0/0)  
0/0

FS CPI  
 FA AB  
 MC CPI: B04-B04A1; B04-C01; D05-C12; D05-H12  
 ABEQ JP 05503000 W UPAB: 19931116

Polypeptides (I) prodn. comprises construction of an in vitro expression unit (II) (contg. a 5' untranslated region with an RNA polymerase binding sequence, a ribosome binding sequence and a translation initiation region), attaching it to at least one semi-random nucleotide sequence and expressing or replication associated sequences followed by translation of the RNA produced to produce polysomes.

The polysomes are bound to a substance of interest, isolated, disrupted to release the RNA which is recovered and converted to a cDNA which can be expressed as a novel polypeptide.

USE - The method allows the isolation of semi-random DNA or RNA sequences that encode novel polypeptides which can bind specific substances.

ABEQ US 5643768 A UPAB: 19970806

A method for isolating a nucleotide sequence which encodes a polypeptide of interest, comprising

transcribing an expression unit which comprises a 5' untranslated region containing an RNA polymerase binding sequence, a ribosome binding sequence, a translation initiation signal, and one or more semi-random nucleotide sequences to produce a mRNA library;

translating in vitro the mRNA library under conditions which maintain polysomes having polypeptide chains attached thereto; contacting the polysomes to a substance of interest and isolating mRNA from the polysomes that specifically bind to the substance of interest.

Dwg.0/0

ABEQ US 5658754 A UPAB: 19970926

A method for producing novel polypeptides, comprising:

(a) constructing an in vitro expression unit comprising a 5' untranslated region containing an RNA polymerase binding site, a ribosome binding site, and a translation initiation signal, said expression unit being capable of producing mRNA;

(b) attaching one or more semi-random polynucleotides to said expression unit;

(c) transcribing or replicating from the RNA polymerase binding site the polynucleotides associated with the expression unit and semi-random polynucleotides to produce mRNA;

(d) translating said mRNA to produce polysomes under conditions sufficient to maintain said polysomes;

(e) binding said polysomes to a substance of interest;

(f) isolating said polysomes that bind to said substance of interest;

(g) disrupting said isolated polysomes to release mRNA;

(h) recovering said mRNA;

(i) constructing cDNA from said recovered mRNA; and

(j) expressing said cDNA to produce novel polypeptides.

Dwg.0/0

=> d his

(FILE 'HOME' ENTERED AT 09:09:12 ON 22 JUL 2003)  
 SET COST OFF

FILE 'HCAPLUS' ENTERED AT 09:09:23 ON 22 JUL 2003

L1 1 S US20020142387/PN  
 E SEKI E/AU  
 L2 3 S E3,E6  
 E KIGAWA T/AU  
 L3 46 S E3,E4  
 E YOKOYAMA S/AU  
 L4 658 S E3-E5,E51

```

      E KIKAWA T/AU
L5      11 S E3,E7
L6      667 S L2-L5
L7      287 S L6 AND PROTEIN
L8      45 S L7 AND CELL(L)FREE
L9      2 S L8 AND FUS?
L10     45 S L7 AND (BIOCHEM?(L)METHOD?)/SC,SX
L11     20 S L8 AND L10
      E TRANSLATION/CT
      E E3+ALL
L12     16019 S E1
      E E2+ALL
L13     32323 S E3,E2
L14     16971 S E2+NT
L15     39 S L7 AND L12-L14
L16     24 S L8 AND L15
L17     30 S L11,L16
L18     63 S L6 AND P/DT
L19     10 S L18 AND L17
L20     53 S L18 NOT L19
L21     20 S L17 NOT L19
L22     30 S L19,L21
L23     50 S L8-L11,L15 NOT L16-L22
      SEL DN AN 20 21
L24     2 S L23 AND E1-E6
L25     32 S L22,L24
      E MOLECULAR STRUCTURE/CT
      E E20+ALL
L26     6957 S E2,E1+NT
      E E16+ALL
L27     960080 S E2,E3,E1+NT
      E CRYSTAL STRUCTURE/CT
      E E8+ALL
L28     6204 S E1
      E E2+ALL
L29     3065 S E1+NT
      E E10+ALL
L30     325985 S E1,E2
L31     304433 S L26-L30 AND PROTEIN
      E FUSION/CT
      E E26+ALL
L32     13851 S E2,E3,E1+NT
L33     5794 S L31 AND L32
L34     15 S L33 AND CELL FREE
L35     56 S L33 AND CELL(L)FREE
L36     14689 S PROTEIN#/SC,SX,CW,BI AND CELL FREE
L37     45380 S PROTEIN#/SC,SX,CW,BI AND CELL(L)FREE
L38     203 S PROTEIN#/SC,SX,CW,BI AND CELLFREE?
L39     1761 S L36-L38 AND FUS?
L40     232 S L32 AND L39
L41     1761 S L39,L40
L42     146 S L41 AND (BIOCHEM?(L)METHOD?)/SC,SX
L43     461 S L41 AND 3/SC,SX
L44     58 S L42,L43 AND (GFP OR GREEN FLUORESCEN?)
L45     2 S L43 AND (3D OR (3 OR THREE OR THIRD) ) (D OR DIM OR DIMENSION?
L46     100 S L12-L14 AND L41
L47     2 S L46 AND (3D OR (3 OR THREE OR THIRD) ) (D OR DIM OR DIMENSION?
L48     37 S L43 AND L46
L49     34 S L25,L45,L47
L50     148 S L44,L46,L48 NOT L49
      SEL DN AN 4-7 10 13 51 57 67 73 76 106
L51     12 S L50 AND E1-E36
L52     46 S L49,L51

```

L53 184 S L42,L43 AND ("E" OR ESCH?) ()COLI  
L54 43 S L53 AND TRANSLAT?  
SEL DN AN 2 3 7 8 10 13 14 20  
L55 8 S L54 AND E37-E60  
L56 54 S L53 AND TRANSCRI?  
L57 31 S L56 NOT L54  
L58 50 S L52,L55  
L59 50 S L58 AND L1-L58  
L60 50 S L59 AND (CELL FREE OR FUS? OR SYNTH? OR VECTOR OR EXPRESS? OR

FILE 'HCAPLUS' ENTERED AT 09:58:46 ON 22 JUL 2003

FILE 'WPIX' ENTERED AT 10:59:22 ON 22 JUL 2003

E JP2001-62703/AP, PRN

L61 1 S E3,E4  
L62 3806 S C12N015-62/IC, ICM, ICS  
L63 7778 S C12N015-11/IC, ICM, ICS  
L64 22927 S C12N015-09/IC, ICM, ICS  
L65 210 S L62 AND L63 AND L64  
L66 167 S L65 AND C07K/IC, ICM, ICS  
L67 89 S L65 AND C07K019/IC, ICM, ICS  
L68 146 S L65 AND C12P021/IC, ICM, ICS  
L69 69 S L67 AND L68  
L70 131 S L66 AND L68  
L71 131 S L69,L70  
L72 1 S L71 AND (GFP OR GREEN FLUORES?)/BIX  
L73 1 S L71 AND G01N024/IC, ICM, ICS  
L74 2 S L71 AND G01N023/IC, ICM, ICS  
L75 129 S L71 NOT L72-L74  
L76 5 S L71 AND (CELL(S)FREE)/BIX  
L77 271 S (CELL FREE OR CELLFREE)/BIX AND L62-L64  
L78 69 S L77 AND L63  
L79 19 S L78 AND C12P021/IC, ICM, ICS, ICA, ICI  
SEL DN AN 17 18  
L80 2 S E1-E4  
L81 49 S L78 AND C07K/IC, ICM, ICS, ICA, ICI  
L82 33 S L81 NOT L79  
L83 17 S L78 NOT L79-L82  
L84 125 S L75 NOT L76,L78-L83  
SEL DN AN 22  
L85 1 S L84 AND E5-E7  
L86 9 S L61,L72,L74,L76,L80,L85  
E KIGAWA T/AU  
L87 8 S E3  
E SEKI E/AU  
L88 19 S E3  
E YOKOYAMA S/AU  
L89 340 S E3,E4  
E KIKAWA/AU  
L90 9 S E7  
L91 371 S L87-L90  
L92 8 S L91 AND C07K/IC, ICM, ICS, ICA, ICI  
L93 16 S L91 AND C12N/IC, ICM, ICS, ICA, ICI  
L94 18 S L91 AND G01N/IC, ICM, ICS, ICA, ICI  
L95 9 S L91 AND C12P/IC, ICM, ICS, ICA, ICI  
L96 8 S L91 AND C12Q/IC, ICM, ICS, ICA, ICI  
L97 30 S L92-L96  
SEL DN AN 5 6 7  
L98 3 S L97 AND E1-E7  
L99 12205 S (CELLFREE OR CELL(S)FREE)/BIX  
L100 1925 S L99 AND C07K/IC, ICM, ICS, ICA, ICI  
L101 1277 S L99 AND C12P021/IC, ICM, ICS, ICA, ICI  
L102 2400 S L100,L101

L103 1308 S L102 AND C12N015/IC, ICM, ICS, ICA, ICI  
L104 113 S L103 AND C12N015-11/IC, ICM, ICS, ICA, ICI  
L105 549 S L103 AND C12N015-09/IC, ICM, ICS, ICA, ICI  
L106 47 S L104 AND L105  
SEL DN AN 1 9  
L107 2 S L106 AND E8-E11  
L108 65 SS C12N015-70/IC, ICM, ICS, ICA, ICI AND L102  
L109 65 S L108 AND L103-L105  
L110 4 S L98, L107  
L111 1 S L86 AND L110  
L112 4 S L110, L111  
L113 8 S L86 NOT L112  
L114 6 S L113 AND L99  
SEL DN AN 4  
L115 1 S L114 AND E12-E13  
L116 5 S L112, L115  
L117 2 S L86 NOT L114-L116

FILE 'WPIX' ENTERED AT 11:46:47 ON 22 JUL 2003

FILE 'WPIX' ENTERED AT 11:47:29 ON 22 JUL 2003

FILE 'WPIX' ENTERED AT 11:48:13 ON 22 JUL 2003

=> d his

(FILE 'HOME' ENTERED AT 09:09:12 ON 22 JUL 2003)  
SET COST OFF

FILE 'HCAPLUS' ENTERED AT 09:09:23 ON 22 JUL 2003

L1 1 S US20020142387/PN  
E SEKI E/AU  
L2 3 S E3,E6  
E KIGAWA T/AU  
L3 46 S E3,E4  
E YOKOYAMA S/AU  
L4 658 S E3-E5,E51  
E KIKAWA T/AU  
L5 11 S E3,E7  
L6 667 S L2-L5  
L7 287 S L6 AND PROTEIN  
L8 45 S L7 AND CELL(L) FREE  
L9 2 S L8 AND FUS?  
L10 45 S L7 AND (BIOCHEM?(L)METHOD?)/SC,SX  
L11 20 S L8 AND L10  
E TRANSLATION/CT  
E E3+ALL  
L12 16019 S E1  
E E2+ALL  
L13 32323 S E3,E2  
L14 16971 S E2+NT  
L15 39 S L7 AND L12-L14  
L16 24 S L8 AND L15  
L17 30 S L11,L16  
L18 63 S L6 AND P/DT  
L19 10 S L18 AND L17  
L20 53 S L18 NOT L19  
L21 20 S L17 NOT L19  
L22 30 S L19,L21  
L23 50 S L8-L11,L15 NOT L16-L22  
SEL DN AN 20 21  
L24 2 S L23 AND E1-E6  
L25 32 S L22,L24  
E MOLECULAR STRUCTURE/CT  
E E20+ALL  
L26 6957 S E2,E1+NT  
E E16+ALL  
L27 960080 S E2,E3,E1+NT  
E CRYSTAL STRUCTURE/CT  
E E8+ALL  
L28 6204 S E1  
E E2+ALL  
L29 3065 S E1+NT  
E E10+ALL  
L30 325985 S E1,E2  
L31 304433 S L26-L30 AND PROTEIN  
E FUSION/CT  
E E26+ALL  
L32 13851 S E2,E3,E1+NT  
L33 5794 S L31 AND L32  
L34 15 S L33 AND CELL FREE  
L35 56 S L33 AND CELL(L) FREE  
L36 14689 S PROTEIN#/SC,SX,CW,BI AND CELL FREE  
L37 45380 S PROTEIN#/SC,SX,CW,BI AND CELL(L) FREE  
L38 203 S PROTEIN#/SC,SX,CW,BI AND CELLFREE?  
L39 1761 S L36-L38 AND FUS?  
L40 232 S L32 AND L39

Jan Delaval  
Reference Librarian  
Biotechnology & Chemical Library  
CM1 1E07 - 703-308-4498  
jan.delaval@uspto.gov

L41 1761 S L39,L40  
 L42 146 S L41 AND (BIOCHEM?(L)METHOD?)/SC,SX  
 L43 461 S L41 AND 3/SC,SX  
 L44 58 S L42,L43 AND (GFP OR GREEN FLUORESCEN?)  
 L45 2 S L43 AND (3D OR (3 OR THREE OR THIRD)())(D OR DIM OR DIMENSION?)  
 L46 100 S L12-L14 AND L41  
 L47 2 S L46 AND (3D OR (3 OR THREE OR THIRD)())(D OR DIM OR DIMENSION?)  
 L48 37 S L43 AND L46  
 L49 34 S L25,L45,L47  
 L50 148 S L44,L46,L48 NOT L49  
 SEL DN AN 4-7 10 13 51 57 67 73 76 106  
 L51 12 S L50 AND E1-E36  
 L52 46 S L49,L51  
 L53 184 S L42,L43 AND ("E" OR ESCH?)()COLI  
 L54 43 S L53 AND TRANSLAT?  
 SEL DN AN 2 3 7 8 10 13 14 20  
 L55 8 S L54 AND E37-E60  
 L56 54 S L53 AND TRANSCRI?  
 L57 31 S L56 NOT L54  
 L58 50 S L52,L55  
 L59 50 S L58 AND L1-L58  
 L60 50 S L59 AND (CELL FREE OR FUS? OR SYNTH? OR VECTOR OR EXPRESS? OR

=> fil hcaplus

FILE 'HCAPLUS' ENTERED AT 09:58:46 ON 22 JUL 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 22 Jul 2003 VOL 139 ISS 4

FILE LAST UPDATED: 21 Jul 2003 (20030721/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d l60 all tot

L60 ANSWER 1 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:242431 HCAPLUS

DN 138:253827

TI Method for increasing the **solubility**, **expression** rate and the activity of **proteins** during **cell-free recombinant** production

IN Watzele, Manfred; Schweizer, Regina; Nemetz, Cordula; Steigerwald, Robin; Emrich, Thomas; Zaiss, Katrin; Fernholz, Erhard; Walckhoff, Baerbel; Schoenfeld, Hans Joachim; Offen, Birgit

PA Roche Diagnostics GmbH, Germany; F. Hoffmann-La-Roche AG

SO PCT Int. Appl., 32 pp.

CODEN: PIXXD2

DT Patent

LA German

IC ICM C12N  
 CC 16-1 (Fermentation and Bioindustrial Chemistry)  
 Section cross-reference(s): 3

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003025116	A2	20030327	WO 2002-EP10329	20020914
	W: JP, US				
	RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR				
	DE 10145694	A1	20030403	DE 2001-10145694	20010917
PRAI	DE 2001-10145694	A	20010917		

AB The invention relates to a method for producing a lysate contg. auxiliary **proteins**. According to said method, a strain, which is suitable for obtaining in vitro **translation** lysates is transformed using a **vector** contg. one or more genes that code for one or more auxiliary **proteins**, the auxiliary **proteins** being **expressed** in said strain and the lysate contg. auxiliary **proteins** being obtained from said strains. The invention also relates to a lysate contg. auxiliary **proteins** that can be obtained according to the inventive method, to blends of said lysates and to the use of the lysates and blends in in vitro **translation** systems. Thus, the inclusion of genes for the chaperonins DnaK, GrpE, and DnaK in an **Escherichia coli** strain that **expresses** human telomerase resulted in increased prodn. of **sol. protein** in a **cell-free** system.

ST **Escherichia cell free translation chaperonin recombinant protein**

IT Bioreactors

(CECF or CFCF; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Chaperonins

RL: BCP (Biochemical process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(DnaJ; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Chaperonins

RL: BCP (Biochemical process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(DnaK; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, animal

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(EPO; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Reporter gene

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(GFP; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Chaperonins

RL: BCP (Biochemical process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(GroEL; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Chaperonins

RL: BCP (Biochemical process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)



(GroES; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT **Proteins**

RL: BCP (Biochemical process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(GrpE; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, microbial

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(MAL; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT **Proteins**

RL: BMF (Bioindustrial manufacture); BIOL (Biological study); PREP (Preparation)  
(MBP (maltose-binding **protein**), **fusion** with erythropoietin; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, animal

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(TEL; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT **Transcription, genetic**

**Translation, genetic**

(**cell-free**; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, microbial

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(dnaJ; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, microbial

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(dnaK; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT **Proteins**

RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)  
(**green fluorescent**; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, microbial

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(groEL; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, microbial

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(groES; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT Gene, microbial

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(grpE; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

- IT **Escherichia coli**  
(lysates of; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT Genetic **vectors**  
**Solubility**  
(method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Chaperonins**  
RL: BCP (Biochemical process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Fusion proteins (chimeric proteins)**  
RL: BMF (Bioindustrial manufacture); BIOL (Biological study); PREP (Preparation)  
(method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Gene**  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Plasmid vectors**  
(pIVEX2.3-GFP; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Plasmid vectors**  
(pIVEX2.4-Rhodanase; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Plasmid vectors**  
(pIVEX2.4b-Mal-Epo; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Plasmid vectors**  
(pIVEX2.4bNde-hTERT; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Plasmid vectors**  
(pRDKJG; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Plasmid vectors**  
(pREP4-groESL; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Human**  
(**proteins**, **expression** of; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Proteins**  
RL: BMF (Bioindustrial manufacture); BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(**recombinant**; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)
- IT **Gene, animal**  
RL: BSU (Biological study, unclassified); BIOL (Biological study)

(rho; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT **Proteins**

RL: BCP (Biochemical process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(trigger factors; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

## IT 95076-93-0

RL: BCP (Biochemical process); CAT (Catalyst use); BIOL (Biological study); PROC (Process); USES (Uses)

(bovine mitochondrial; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

## IT 9001-92-7, Protease 9001-99-4, RNase 37228-74-3, Exonuclease

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(**cells** that have low levels or are deficient in; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT 11096-26-7DP, Erythropoietin, **fusion protein** with maltose-binding **protein** 120178-12-3P, Telomerase

RL: BMF (Bioindustrial manufacture); BIOL (Biological study); PREP (Preparation)

(human; method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

## IT 9026-04-4P, Rhodanase

RL: BMF (Bioindustrial manufacture); BIOL (Biological study); PREP (Preparation)

(method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

IT 37318-49-3, **Protein** disulfide isomerase

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(method for increasing **soly.**, **expression** rate and activity of **proteins** during **cell-free recombinant** prodn.)

L60 ANSWER 2 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:187867 HCAPLUS

DN 138:201358

TI Nucleic acid ligation via primer-less PCR for **fusion protein expression** in **cell-free** translation

IN Nemoto, Naoto

PA Mitsubishi Chemical Corp., Japan

SO Jpn. Kokai Tokkyo Koho, 16 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM C12N015-09

ICS C12P021-02

CC 9-15 (Biochemical Methods)

Section cross-reference(s): 3, 16

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2003070483	A2	20030311	JP 2001-270323	20010906
PRAI	JP 2001-270323		20010906		

AB A method for ligating nucleic acid via primer-less PCR using a pair of complementary and overlapping single-stranded or double-stranded DNA, is

disclosed. One of the DNA codes for a desired **protein** and the other codes for an .alpha.-helix contg. globular **protein** as support for **expressing** or displaying **proteins** or peptides as **fusion protein** in **cell-free** translation system. DNA polymerase, Taq polymerase, in particular, is used for PCR and RNA polymerase is used for transcription. DNase is used to degrade contaminating DNA. Nucleic acid derivs. such as puromycin, 3-N-aminoacylpuromycin aminonucleoside (PANS-amino acid), or 3'-N-aminoacyl adenosine aminonucleosides (AANS-amino acid), are linked to **fusion protein** coding mRNA via a linker (spacer). Polyethylene or polyethylene glycol may be used as spacer. A mutated POU-specific **domain** of transcription factor Oct-1 having a Cys residue replaced by Ala, was used as support for **expression** of peptide library in **cell-free** translation system. Carbohydrate-binding peptides, binding N-acetylglucosamine in the presence of calcium was screened from the peptide library.

ST nucleic acid ligation primer less PCR; **fusion protein expression cell free** translation

IT Nucleoside analogs

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(3-N-aminoacylpuromycin amino, as nucleic acid deriv.; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free** translation)

IT Nucleoside analogs

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(3'-N-amino acyl adenosine amino, as nucleic acid deriv.; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free** translation)

IT Transcription factors

RL: BUU (Biological use, unclassified); PRP (Properties); BIOL (Biological study); USES (Uses)

(NF-III (nuclear factor III), POU-specific **domain** of; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free** translation)

IT Nucleic acids

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(analog; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free** translation)

IT Translation, genetic

(**cell-free**; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free** translation)

IT DNA

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(double-stranded; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free** translation)

IT Proteins

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(globular; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free** translation)

IT DNA

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(linker; nucleic acid ligation via primer-less PCR for **fusion protein expression in cell-free**

- translation)
- IT PCR (polymerase chain reaction)  
Peptide library  
**Protein** sequences  
Transcription, genetic  
(nucleic acid ligation via primer-less PCR for **fusion**  
**protein expression in cell-free**  
translation)
- IT **Fusion proteins (chimeric proteins)**  
Peptides, preparation  
**Proteins**  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP  
(Preparation)  
(nucleic acid ligation via primer-less PCR for **fusion**  
**protein expression in cell-free**  
translation)
- IT Nucleic acids  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(nucleic acid ligation via primer-less PCR for **fusion**  
**protein expression in cell-free**  
translation)
- IT Polyoxyalkylenes, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(nucleic acid ligation via primer-less PCR for **fusion**  
**protein expression in cell-free**  
translation)
- IT DNA  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(single-stranded; nucleic acid ligation via primer-less PCR for  
**fusion protein expression in cell**  
**-free** translation)
- IT **Proteins**  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(.alpha.-helix-contg.; nucleic acid ligation via primer-less PCR for  
**fusion protein expression in cell**  
**-free** translation)
- IT 53-79-2, Puromycin 53-79-2D, Puromycin, 3'-N-aminoacyl aminonucleoside  
58-61-7D, Adenosine, 3'-N-aminoacyl aminonucleosides  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(as nucleic acid deriv.; nucleic acid ligation via primer-less PCR for  
**fusion protein expression in cell**  
**-free** translation)
- IT 9003-98-9, DNase  
RL: BUU (Biological use, unclassified); CAT (Catalyst use); BIOL  
(Biological study); USES (Uses)  
(for degrading contaminating DNA; nucleic acid ligation via primer-less  
PCR for **fusion protein expression in**  
**cell-free** translation)
- IT 25322-68-3, Polyethylene glycol  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(nucleic acid ligation via primer-less PCR for **fusion**  
**protein expression in cell-free**  
translation)
- IT 500256-17-7 500256-18-8 500256-19-9 500256-20-2 500256-21-3  
500256-22-4 500256-23-5 500256-24-6  
RL: PRP (Properties)  
(unclaimed nucleotide sequence; nucleic acid ligation via primer-less

PCR for **fusion protein expression** in **cell-free** translation)

IT 500256-25-7  
 RL: PRP (Properties)  
 (unclaimed **protein** sequence; nucleic acid ligation via primer-less PCR for **fusion protein expression** in **cell-free** translation)

IT 9002-88-4, Polyethylene  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (use as linker; nucleic acid ligation via primer-less PCR for **fusion protein expression** in **cell-free** translation)

IT 9012-90-2, DNA polymerase  
 RL: BUU (Biological use, unclassified); CAT (Catalyst use); BIOL (Biological study); USES (Uses)  
 (use for PCR; nucleic acid ligation via primer-less PCR for **fusion protein expression** in **cell-free** translation)

IT 9014-24-8, RNA polymerase  
 RL: BUU (Biological use, unclassified); CAT (Catalyst use); BIOL (Biological study); USES (Uses)  
 (use for transcription; nucleic acid ligation via primer-less PCR for **fusion protein expression** in **cell-free** translation)

L60 ANSWER 3 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:187866 HCAPLUS

DN 138:201357

TI Globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system

IN Nemoto, Naoto

PA Mitsubishi Chemical Corp., Japan

SO Jpn. Kokai Tokkyo Koho, 15 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM C12N015-09

ICS C07K014-00; C07K019-00; C12P021-02

CC 9-15 (Biochemical Methods)

Section cross-reference(s): 3, 16

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2003070482	A2	20030311	JP 2001-270322	20010906
PRAI	JP 2001-270322		20010906		

AB. Use of .alpha.-helix contg. globular **protein** as support for **expressing** or displaying **proteins** or peptides as **fusion protein** in **cell-free** translation system, is disclosed. Nucleic acid derivs. such as puromycin, 3-N-aminoacylpuromycin aminonucleoside (PANS-amino acid), or 3'-N-aminoacyl adenosine aminonucleosides (AANS-amino acid), are linked to **fusion protein** coding mRNA via a linker (spacer). Polyethylene or polyethylene glycol may be used as spacer. A mutated POU-specific **domain** of transcription factor Oct-1 having a Cys residue replaced by Ala, was used as support for **expression** of peptide library in **cell-free** translation system. Carbohydrate-binding peptides, binding N-acetylglucosamine in the presence of calcium was screened from the peptide library.

ST globular **protein** support displaying **fusion protein cell free** translation

IT Nucleoside analogs

- RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(3'-N-aminoacylpuromycin amino, as nucleic acid deriv.; globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT Nucleoside analogs  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(3'-N-amino acyl adenosine amino, as nucleic acid deriv.; globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT Transcription factors  
RL: BUU (Biological use, unclassified); PRP (Properties); BIOL (Biological study); USES (Uses)  
(NF-III (nuclear factor III), POU-specific **domain** of; globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT Nucleic acids  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(analog; globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT **Translation, genetic**  
(**cell-free**; globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT Peptide library  
**Protein** sequences  
(globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT **Fusion proteins (chimeric proteins)**  
Peptides, preparation  
**Proteins**  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT Polyoxyalkylenes, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT **Proteins**  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(globular; globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT **Proteins**  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(.alpha.-helix-contg.; globular **protein** as support for **expressing** or displaying **fusion proteins** in **cell-free** translation system)
- IT 500252-38-0  
RL: BUU (Biological use, unclassified); PRP (Properties); BIOL (Biological

study); USES (Uses)  
 (amino acid sequence; globular **protein** as support for  
**expressing** or displaying **fusion proteins** in  
**cell-free** translation system)

IT 53-79-2, Puromycin 53-79-2D, Puromycin, 3-N-aminoacyl aminonucleoside  
 58-61-7D, Adenosine, 3'-N-aminoacyl aminonucleosides  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (as nucleic acid deriv.; globular **protein** as support for  
**expressing** or displaying **fusion proteins** in  
**cell-free** translation system)

IT 25322-68-3, Polyethylene glycol  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (globular **protein** as support for **expressing** or  
 displaying **fusion proteins** in **cell-**  
**free** translation system)

IT 500256-26-8 500256-27-9 500256-28-0 500256-29-1 500256-30-4  
 500256-31-5 500256-32-6 500256-33-7  
 RL: PRP (Properties)  
 (unclaimed nucleotide sequence; globular **protein** as support  
 for **expressing** or displaying **fusion**  
**proteins** in **cell-free** translation system)

IT 9002-88-4, Polyethylene  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (use as linker; globular **protein** as support for  
**expressing** or displaying **fusion proteins** in  
**cell-free** translation system)

L60 ANSWER 4 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
 AN 2003:150699 HCAPLUS  
 DN 138:199905  
 TI Method for determining gene **expression** in a **cell-**  
**free transcription/translation** system  
 IN Erdmann, Volker A.; Lamla, Thorsten; Stiege, Wolfgang  
 PA Germany  
 SO Ger. Offen., 8 pp.  
 CODEN: GWXXBX  
 DT Patent  
 LA German  
 IC ICM C12N015-00  
 ICS C12M001-42; C12P021-00  
 CC 3-1 (Biochemical Genetics)  
 Section cross-reference(s): 9

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 10137792	A1	20030227	DE 2001-10137792	20010806
	WO 2003016526	A1	20030227	WO 2002-DE2672	20020717
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
	CO, CR, CU, CZ, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM,				
	HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,				
	LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT,				
	RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG,				
	US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AT, BE, BG,				
	CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL,				
	PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,				
	NE, SN, TD, TG				

PRAI DE 2001-10137792 A 20010806

AB The invention relates to a method using a flow reaction chamber for  
 studying gene **expression** in a **cell-free**



**transcription/translation** system. The reaction soln. contains all necessary components of the **transcription/translation** app., amino acids, nucleotides, energy supplying and metab. components necessary for the **synthesis**. During **translation** the produced **proteins** are immobilized on a matrix.

- ST gene **expression** detection **cell free**  
**transcription translation** system
- IT **Escherichia coli**  
(S30 D10, lysate of; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Pumps  
(centrifugal, for maintaining the reaction flow in the reactor; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Amino acids, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(component of reaction mix; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Gene  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(**expression**; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT UV and visible spectroscopy  
(for detg. **protein** concn.; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Matrix media  
(for specific **protein** binding, hydrophilic/hydrophobic, antigen/antibody, metal ion; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Antibodies  
Antigens  
RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(for specific **protein** binding; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT **Transcription, genetic**  
**Translation, genetic**  
(method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT **Fusion proteins (chimeric proteins)**  
RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Immobilization, molecular  
(of **protein** products; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Reactors  
(reaction chamber of **transcription/translation** system; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT Membranes, nonbiological  
(semipermeable, component of the reactor; method for detg. gene **expression** in a **cell-free transcription/translation** system)

- IT 147395-23-1 205938-74-5  
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
(amino acid sequence; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT 50812-37-8, Glutathione-S-transferase  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(as **protein** N- or C-terminal **fusion** partner; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT 56-65-5, ATP, biological studies 63-39-8, UTP 65-47-4, CTP 86-01-1, GTP 2466-09-3, Diphosphoric acid  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(component of reaction mix; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT 9013-20-1, Streptavidin  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(for binding WSHPQFEK and AWRHPQFGG amino acid sequences; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT 71-00-1, Histidine, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(frequency of in **protein** N-terminal or C-terminal; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT 15158-11-9, Cu 2+, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(matrix component; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT 1306-06-5, Hydroxyapatite  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(matrix; method for detg. gene **expression** in a **cell-free transcription/translation** system)
- IT 14701-22-5, Nickel 2+, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(method for detg. gene **expression** in a **cell-free transcription/translation** system)

RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Anon; Nucl Acids Red 2001, V29(15), Pe73

L60 ANSWER 5 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:133583 HCAPLUS

DN 138:183528

TI Method for detecting interaction between substance and **protein** using **cell-free** translation system, method for screening **protein** interacting with substance, and method for forming complex of substance and **protein** interacting with the substance

IN Yanagawa, Hiroshi; Miyamoto, Etsuko; Matsumura, Nobutaka; Doi, Nobuhide; Tateyama, Seiji; Ishizaka, Masamichi; Horisawa, Kenichi

PA Keio University, Japan

SO PCT Int. Appl., 114 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

IC ICM G01N033-50  
ICS G01N033-15; G01N033-566  
CC 9-16 (Biochemical Methods)  
Section cross-reference(s): 3, 6  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003014734	A1	20030220	WO 2002-JP8078	20020807
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
PRAI	JP 2001-239175	A	20010807		
	JP 2002-172129	A	20020612		
AB	<p>A method for detecting an interaction between a substance (bait) and a <b>protein</b> (prey) using a <b>cell-free</b> translation system is provided, in which the bait is brought contact with the prey, and the complex formed by the contact is detected. In this method, a <b>fusion protein</b> of the prey with a <b>protein</b> usable as a label for detection is designed while the bait is added with a modification for sepn. Then, mRNA encoding the <b>fusion protein</b> contg. the prey is translated in a <b>cell-free</b> translation system in the presence of the bait, and thereby, the translation step is coupled with the interaction step. Thus, the prey is formed in the <b>cell-free</b> translation system, and then, brought into contact with the bait. The complex between the bait and the prey is detected with the use of the modification for sepn. and the label for detection. Thus, the anal. of an interaction between <b>proteins</b>, between a nucleic acid and a <b>protein</b>, or else is performed in vitro at a high sensitivity. In case where the bait is a <b>protein</b>, alternatively, mRNA encoding the <b>fusion protein</b> contg. the bait and mRNA encoding the <b>protein</b> contg. the prey or its <b>fusion protein</b> are translated in a <b>cell-free</b> cotranslation system, and thereby, the translation are coupled with the interaction step. Thus, the bait and prey formed are allowed to interact with each other, and the complex is detected with the use of the modification for sepn. and the label for detection. Thereby, the anal. of an interaction between <b>proteins</b>, between a nucleic acid and a <b>protein</b>, or else is performed in vitro at a high sensitivity.</p>				
ST	<b>protein</b> interaction <b>cell free</b> translation system; in vivo virus STABLE <b>protein</b> DNA interaction				
IT	<b>Proteins</b>				
	RL: BSU (Biological study, unclassified); BIOL (Biological study) (DNA-binding; methods for detecting <b>protein-protein</b> or <b>protein-nucleic acid</b> interaction using <b>cell-free</b> translation system)				
IT	<b>Translation, genetic</b> ( <b>cell-free</b> ; methods for detecting <b>protein-protein</b> or <b>protein-nucleic acid</b> interaction using <b>cell-free</b> translation system)				
IT	Nucleic acid library (dT priming cDNA; methods for detecting <b>protein-protein</b> or <b>protein-nucleic acid</b> interaction using <b>cell-free</b> translation system)				
IT	Molecular association				

(methods for detecting **protein-protein** or **protein-nucleic acid** interaction using **cell-free** translation system)

IT Avidins  
    **Fusion proteins (chimeric proteins)**  
Nucleic acids  
    **Proteins**  
Transcription factors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
    (methods for detecting **protein-protein** or **protein-nucleic acid** interaction using **cell-free** translation system)

IT mRNA  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
    (methods for detecting **protein-protein** or **protein-nucleic acid** interaction using **cell-free** translation system)

IT Nucleic acid library  
    (random priming cDNA; methods for detecting **protein-protein** or **protein-nucleic acid** interaction using **cell-free** translation system)

IT Recombination, genetic  
    (translocation; methods for detecting **protein-protein** or **protein-nucleic acid** interaction using **cell-free** translation system)

IT 53-79-2, Puromycin 58-85-5, Biotin 9013-20-1, Streptavidin  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
    (methods for detecting **protein-protein** or **protein-nucleic acid** interaction using **cell-free** translation system)

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Keio University; WO 0248347 A 2002 HCAPLUS  
(2) Keio University; JP 2002176987 A 2002 HCAPLUS  
(3) Mitsubishi Chemical Corp; US 6361943 A 1998 HCAPLUS  
(4) Mitsubishi Chemical Corp; EP 962527 A 1998 HCAPLUS  
(5) Mitsubishi Chemical Corp; WO 9816636 A 1998 HCAPLUS  
(6) Mitsubishi Chemical Corp; JP 11-322781 A 1999 HCAPLUS  
(7) Mitsubishi Chemical Corp; JP 2000139468 A 2000 HCAPLUS  
(8) Mitsubishi Chemical Corp; JP 2001128690 A 2001 HCAPLUS  
(9) The Institute Of Physical And Chemical Research; WO 0068690 A 2001 HCAPLUS  
(10) The Institute Of Physical And Chemical Research; EP 1182458 A 2001 HCAPLUS  
(11) The Institute Of Physical And Chemical Research; JP 2001027633 A 2001 HCAPLUS

L60 ANSWER 6 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2003:133461 HCAPLUS  
DN 138:182052  
TI Engineered **Escherichia coli** tyrosyl-tRNA  
    **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system  
IN Kiga, Daisuke; Sakamoto, Kensaku; Hirao, Ichiro; Yokoyama, Shigeyuki  
PA Japan Science and Technology Corporation, Japan  
SO PCT Int. Appl., 49 pp.  
CODEN: PIXXD2  
DT **Patent**  
LA Japanese  
IC ICM C12N015-52  
ICS C12N009-00; C07B061-00; G01N033-48; G06F017-30; G06F017-50  
CC 3-2 (Biochemical Genetics)

Section cross-reference(s): 7, 9, 16

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003014354	A1	20030220	WO 2002-JP118	20020111
	W: CA, JP, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				

PRAI JP 2001-234135 A 20010801

AB Genetically modified tyrosyl-tRNA **synthases** more efficiently incorporating unnatural amino acids, particularly 3-substituted tyrosine analogs, are provided. A method of designing such tyrosyl-tRNA **synthase** based on the three-dimensional structure anal. of the complex of tyrosyl-tRNA **synthase** and tyrosyl-AMP is also provided. Use of such modified tyrosyl-tRNA **synthases** for site-specific incorporation of an unnatural amino acid into **proteins** is also provided. Tyrosyl-tRNA **synthetase** (TyrRS) from **Escherichia coli** was engineered to preferentially recognize 3-iodo-L-tyrosine rather than L-tyrosine for the site-specific incorporation of 3-iodo-L-tyrosine into **proteins** in eukaryotic translation systems. The wild-type TyrRS does not recognize 3-iodo-L-tyrosine, because of the bulky iodine substitution. On the basis of the reported crystal structure of Bacillus stearothermophilus TyrRS, three residues, Y37, Q179, and Q195, in the L-tyrosine-binding site were chosen for mutagenesis. Thirty-four single amino acid replacements and 16 of their combinations were screened by in vitro biochem. assays. A combination of the Y37V and Q195C mutations changed the amino acid specificity in such a way that the variant TyrRS activates 3-iodo-L-tyrosine 10-fold more efficiently than L-tyrosine. This engineered **enzyme**, TyrRS(V37C195), was tested for use in the wheat germ **cell-free** translation system, which has recently been significantly improved, and is now as productive as conventional **recombinant** systems. During the translation in the wheat germ system, an **E. coli** suppressor tRNA<sup>Tyr</sup> was not aminoacylated by the wheat germ **enzymes**, but was aminoacylated by the **E. coli** TyrRS(V37C195) variant with 3-iodo-L-tyrosine. After the use of the 3-iodotyrosyl-tRNA in translation, the resultant uncharged tRNA could be aminoacylated again in the system. A mass spectrometric anal. of the produced **protein** revealed that more than 95% of the amino acids incorporated for an amber codon were iodotyrosine, whose concn. was only twice that of L-tyrosine in the translation. Therefore, the variant **enzyme**, 3-iodo-L-tyrosine, and the suppressor tRNA can serve as an addnl. set orthogonal to the 20 endogenous sets in eukaryotic in vitro translation systems.

ST engineering Escherichia tyrosyl tRNA **synthetase** unnatural amino acid translation; tyrosyl tRNA **synthetase** substrate specificity change genetic engineering; iodotyrosine incorporation tyrosyl tRNA **synthetase** Escherichia mutagenesis

IT Codons

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(amber, use in unnatural amino acid incorporation; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)

IT Translation, genetic

(**cell-free**; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)

IT Aminoacylation

Genetic engineering

**Protein engineering**

(engineered **Escherichia coli** tyrosyl-tRNA  
**synthetase** for site-specific incorporation of an unnatural  
amino acid into **proteins** in **cell-free**  
translation system)

IT **Proteins**

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP  
(Preparation)

(incorporating unnatural amino acids; engineered **Escherichia**  
**coli** tyrosyl-tRNA **synthetase** for site-specific  
incorporation of an unnatural amino acid into **proteins** in  
**cell-free** translation system)

IT **Ras proteins**

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP  
(Preparation)

(p21c-Ha-ras, incorporating unnatural amino acids; engineered  
**Escherichia coli** tyrosyl-tRNA **synthetase**  
for site-specific incorporation of an unnatural amino acid into  
**proteins** in **cell-free** translation system)

IT **Conformation**

(**protein**, of TyrRS-tyrosyl-AMP complex, **protein**.  
engineering based on; engineered **Escherichia coli**  
tyrosyl-tRNA **synthetase** for site-specific incorporation of an  
unnatural amino acid into **proteins** in **cell-**  
**free** translation system)

IT **Mutagenesis**

(site-directed, substitution; engineered **Escherichia**  
**coli** tyrosyl-tRNA **synthetase** for site-specific  
incorporation of an unnatural amino acid into **proteins** in  
**cell-free** translation system)

IT **Enzyme functional sites**

(substrate-binding; engineered **Escherichia coli**  
tyrosyl-tRNA **synthetase** for site-specific incorporation of an  
unnatural amino acid into **proteins** in **cell-**  
**free** translation system)

IT **tRNA**

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)

(suppressor, use in unnatural amino acid incorporation; engineered  
**Escherichia coli** tyrosyl-tRNA **synthetase**  
for site-specific incorporation of an unnatural amino acid into  
**proteins** in **cell-free** translation system)

IT **Eukaryota****Prokaryote**

(transformed with modified tyrosyl-tRNA **synthase** gene;  
engineered **Escherichia coli** tyrosyl-tRNA  
**synthetase** for site-specific incorporation of an unnatural  
amino acid into **proteins** in **cell-free**  
translation system)

IT **Escherichia coli**

(tyrosyl-tRNA **synthetase** from; engineered **Escherichia**  
**coli** tyrosyl-tRNA **synthetase** for site-specific  
incorporation of an unnatural amino acid into **proteins** in  
**cell-free** translation system)

IT 234771-42-7D, Tyrosyl-AMP, complex with tyrosyl-tRNA **synthetase**

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(3D structure of; engineered **Escherichia coli**  
tyrosyl-tRNA **synthetase** for site-specific incorporation of an  
unnatural amino acid into **proteins** in **cell-**  
**free** translation system)

## IT 52-90-4, L-Cysteine, biological studies 56-45-1, L-Serine, biological studies 70-47-3, L-Asparagine, biological studies

RL: BSU (Biological study, unclassified); BIOL (Biological study)

- (at position 195 in mutant; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)
- IT 56-85-9, L-Glutamine, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (at position 195 in wild type; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)
- IT 56-41-7, L-Alanine, biological studies 61-90-5, L-Leucine, biological studies 72-18-4, L-Valine, biological studies 73-32-5, L-Isoleucine, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (at position 37 in mutant; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)
- IT 60-18-4, L-Tyrosine, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (at position 37 in wild type; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)
- IT 9023-45-4, Tyrosyl-tRNA **synthetase**  
 RL: BSU (Biological study, unclassified); CAT (Catalyst use); PRP (Properties); BIOL (Biological study); USES (Uses)  
 (engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)
- IT 60-18-4D, L-Tyrosine, 3-halo-substituted analogs 60-18-4D, L-Tyrosine, 3-substituted analogs 70-78-0, 3-Iodo-L-tyrosine  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (incorporation of; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)
- IT 497985-48-5 497985-49-6 497985-50-9 497985-51-0 497985-52-1  
 497985-53-2 497985-54-3 497985-55-4 497985-56-5 497985-57-6  
 497985-58-7  
 RL: PRP (Properties)  
 (unclaimed nucleotide sequence; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)
- IT 497985-47-4 497985-59-8  
 RL: PRP (Properties)  
 (unclaimed **protein** sequence; engineered **Escherichia coli** tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in **cell-free** translation system)

RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE

- (1) Bain, J; Biochemistry 1991, V30(22), P5411 HCAPLUS
- (2) Barker, D; FEBS Letters 1982, V150(2), P419 HCAPLUS
- (3) Brick, P; J Mol Biol 1989, V208(1), P83 HCAPLUS
- (4) Hamano-Takaku, F; J Biol Chem 2000, V275(51), P40324 HCAPLUS
- (5) Ohno, S; J Biochem 2001, V130(3), P417 HCAPLUS
- (6) Winter, G; Eur J Biochem 1983, V132(2), P383 HCAPLUS

L60 ANSWER 7 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
 AN 2003:132002 HCAPLUS

TI Present status of **cell-free protein synthesis**  
 AU **Kigawa, Takanori; Yokoyama, Shigeyuki**  
 CS Institute of Physical and Chemical Research, Japan  
 SO Posutoshikuensu Tanpakushitsu Jikkenho (2002), Volume 2, 14-24.  
 Editor(s): Oshima, Tairo. Publisher: Tokyo Kagaku Dojin, Tokyo, Japan.  
 CODEN: 69DQE2  
 DT Conference  
 LA Japanese  
 CC **9 (Biochemical Methods)**  
 AB Unavailable

L60 ANSWER 8 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
 AN 2003:53294 HCAPLUS  
 DN 138:299270  
 TI **Protein expression** systems for structural genomics and proteomics  
 AU **Yokoyama, Shigeyuki**  
 CS RIKEN Genomic Sciences Center, 1-7-22 Suehiro-cho, Tsurumi, Yokohama, 230-0045, Japan  
 SO Current Opinion in Chemical Biology (2003), 7(1), 39-43  
 CODEN: COCBF4; ISSN: 1367-5931  
 PB Elsevier Science Ltd.  
 DT Journal; General Review  
 LA English  
 CC 6-0 (General Biochemistry)  
 Section cross-reference(s): 3  
 AB A review. One of the key steps of structural genomics and proteomics is high-throughput **expression** of many target **proteins**. Gene **cloning**, esp. by ligation-independent **cloning** techniques, and **recombinant protein expression** using microbial hosts such as **Escherichia coli** and the yeast *Pichia pastoris* are well optimized and further robotized. **Cell-free protein synthesis** systems have been developed for large-scale prodn. of **protein** samples for NMR (stable-isotope labeling) and X-ray crystallog. (selenomethionine substitution). **Protein folding** is still a major bottleneck in **protein expression**. **Cell**-based and **cell-free** methods for screening of suitable samples for structure detn. have been developed for achieving a high success rate.  
 ST review **protein** sample **expression** system structural genomics proteomics  
 IT **Translation, genetic**  
 (cell-free, for **protein** sample prepn.; **protein expression** systems for structural genomics and proteomics)  
 IT High throughput screening  
 (**protein expression** systems for structural genomics and proteomics)  
 IT **Proteins**  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (**protein expression** systems for structural genomics and proteomics)  
 IT Sample preparation  
 (**protein** sample prepn.; **protein expression** systems for structural genomics and proteomics)  
 IT **Escherichia coli**  
*Komagataella pastoris*  
 (**protein** sample prodn. in; **protein expression** systems for structural genomics and proteomics)  
 IT Genome



(structural genomics, **protein** sample prepn.; **protein expression** systems for structural genomics and proteomics)

IT Proteome

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(structural proteomics, **protein** sample prepn.;  
**protein expression** systems for structural genomics  
and proteomics)

RE.CNT 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Boettner, M; J Biotechnol 2002, V99, P51 HCAPLUS
- (2) Christendat, D; Nat Struct Biol 2000, V7, P903 HCAPLUS
- (3) Christendat, D; Prog Biophys Mol Biol 2000, V73, P339 HCAPLUS
- (4) de Lamotte, F; Protein Expr Purif 2001, V22, P318 HCAPLUS
- (5) Dieckman, L; Protein Expr Purif 2002, V25, P1 HCAPLUS
- (6) Ding, H; Acta Crystallogr D Biol Crystallogr 2002, V58, P2102
- (7) Dyson, H; Curr Opin Struct Biol 2002, V12, P54 HCAPLUS
- (8) Edwards, A; Nat Struct Biol 2000, V7(Suppl), P970
- (9) Gilbert, M; Curr Opin Chem Biol 2002, V6, P102 HCAPLUS
- (10) Guignard, L; FEBS Lett 2002, V524, P159 HCAPLUS
- (11) Hendrickson, W; EMBO J 1990, V9, P1665 HCAPLUS
- (12) Hurley, J; Trends Biochem Sci 2002, V27, P48 HCAPLUS
- (13) Jiang, X; FEBS Lett 2002, V514, P290 HCAPLUS
- (14) Kawasaki, M; Biochem Biophys Res Commun 2001, V280, P842 HCAPLUS
- (15) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (16) Kigawa, T; J Biomol NMR 1995, V6, P129 HCAPLUS
- (17) Kigawa, T; J Struct Funct Genomics 2002, V2, P29 HCAPLUS
- (18) Kim, S; Curr Opin Struct Biol 2000, V10, P380 HCAPLUS
- (19) Knaust, R; Anal Biochem 2001, V297, P79 HCAPLUS
- (20) Lamla, T; Mol Cell Proteomics 2002, V1, P466 HCAPLUS
- (21) Lesley, S; Proc Natl Acad Sci USA 2002, V99, P11664 HCAPLUS
- (22) Madin, K; Proc Natl Acad Sci USA 2000, V97, P559 HCAPLUS
- (23) Mittl, P; Curr Opin Chem Biol 2001, V5, P402 HCAPLUS
- (24) Pedelacq, J; Nat Biotechnol 2002, V20, P927 HCAPLUS
- (25) Sawasaki, T; FEBS Lett 2002, V514, P102 HCAPLUS
- (26) Sawasaki, T; Proc Natl Acad Sci USA 2002, V99, P14652 HCAPLUS
- (27) Shimizu, Y; Nat Biotechnol 2001, V19, P751 HCAPLUS
- (28) Spirin, A; Science 1988, V242, P1162 HCAPLUS
- (29) Stols, L; Protein Expr Purif 2002, V25, P8 HCAPLUS
- (30) Tabuchi, M; Proteomics 2002, V2, P430 HCAPLUS
- (31) Vitkup, D; Nat Struct Biol 2001, V8, P559 HCAPLUS
- (32) Waldo, G; Nat Biotechnol 1999, V17, P691 HCAPLUS
- (33) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS
- (34) Yee, A; Proc Natl Acad Sci USA 2002, V99, P1825 HCAPLUS
- (35) Yokoyama, S; Nat Struct Biol 2000, V7(Suppl), P943
- (36) Zhou, P; J Biomol NMR 2001, V20, P11 HCAPLUS

L60 ANSWER 9 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:39992 HCAPLUS

DN 138:286030

TI **Protein synthesis** and co-translational **folding**  
in **cell-free** translation systems

AU Spirin, Alexander S.

CS Inst. of Protein Res., Russian Acad. of Sci, Pushchino, 142292, Russia

SO Recombinant Protein Production with Prokaryotic and Eukaryotic Cells: A  
Comparative View on Host Physiology, Selected Articles from the Meeting of  
the EFB Section on Microbial Physiology, Semmering, Austria, Oct. 5-8,  
2000 (2001), Meeting Date 2000, 1-15. Editor(s): Merten, Otto-Wilhelm.  
Publisher: Kluwer Academic Publishers, Dordrecht, Neth.  
CODEN: 69DLJQ; ISBN: 0-7923-7137-2

DT Conference; General Review

LA English

CC 16-0 (Fermentation and Bioindustrial Chemistry)

AB A review, with refs. The **cell-free** methodol. for the

**synthesis** of **functionally** active **proteins** is considered, and the so-called continuous **cell-free** translation and transcription-translation systems are described. The continuous **cell-free** systems for gene **expression** are based on the use of a porous barrier that retains the high-mol.-wt. components of the **protein-synthesizing** machinery within a defined reaction compartment, and at the same time provides the continuous feeding with substrates (NTPs and amino acids) and the removal of reaction products. There are two versions of the continuous systems: the flow version (continuous-flow **cell-free**, or CFCF systems) and the dialysis version (continuous-exchange **cell-free**, or CECF systems). Both versions have been shown to provide a prolonged **synthesis** of **proteins**, as compared with std. (batch) **cell-free** systems, and correspondingly a significantly higher yield of **proteins synthesized**. The **synthesis** of **fusion proteins** and the direct **expression** of PCR products in **cell-free** systems are discussed as promising methodol. approaches in a no. of cases. Using the monitoring of polypeptide elongation in **cell-free** systems the evidence is presented that the **folding** of **synthesized** polypeptides into **functional protein** globules proceeds on ribosomes during translation (co-translational **protein folding**).

ST review **protein synthesis folding**  
**cell free** translation system

IT **Translation, genetic**  
 (cell-free; **protein synthesis**  
 and co-translational **folding** in **cell-free**  
 translation systems)

IT Biotechnology  
**Protein folding**  
 (**protein synthesis** and co-translational  
**folding** in **cell-free** translation systems)

IT **Proteins**  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (**protein synthesis** and co-translational  
**folding** in **cell-free** translation systems)

RE.CNT 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE

- (1) Alakhov, Y; US 5478730 1995 HCAPLUS
- (2) Alexandrov, A; Biochem Mol Biol Intern 1996, V38, P1111 HCAPLUS
- (3) Baranov, V; Gene 1989, V84, P463 HCAPLUS
- (4) Baranov, V; Methods in Enzymology, Recombinant DNA, Part H 1993, V217, P123 HCAPLUS
- (5) Birjukov, S; WO 0058493 A 1999 HCAPLUS
- (6) Chekulayeva, M; Biochem Biophys Res Commun, in press 2001
- (7) Davis, J; Promega Notes Magazine 1996, 56, P14
- (8) DeVries, J; Proc Natl Acad Sci USA 1967, V57, P1011
- (9) Endo, Y; J Biotech 1992, V25, P221 HCAPLUS
- (10) Endo, Y; Tokishima J Exp Med 1993, V40(1-2), P13 MEDLINE
- (11) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (12) Kigawa, T; J Biochem (Japan) 1991, V110, P166 HCAPLUS
- (13) Kim, D; Biotech Bioengineering 1999, V66, P180 HCAPLUS
- (14) Kim, D; Biotech Prog 2000, V16, P385 HCAPLUS
- (15) Kim, D; Biotechnol Prog 1996, V12, P645 HCAPLUS
- (16) Kolb, V; Biotech Lett 1996, V18, P1447 HCAPLUS
- (17) Kolb, V; EMBO J 1994, V13, P3631 HCAPLUS
- (18) Kolb, V; J Biol Chem 2000, V275, P16597 HCAPLUS
- (19) Kolosov, M; Biotech Appl Biochem 1992, V16, P125 HCAPLUS
- (20) Komar, A; FEBS Lett 1993, V326, P261 HCAPLUS
- (21) Komar, A; J Biol Chem 1997, V272, P10646 HCAPLUS
- (22) Kudlicki, W; Anal Biochem 1992, V206, P389 HCAPLUS

- (23) Madin, K; Proc Natl Acad Sci USA 2000, V97, P559 HCAPLUS
- (24) Martemyanov, K; Biotech Lett 1996, V18, P1357 HCAPLUS
- (25) Martemyanov, K; Doklady Biochemistry (Russia) 1997, V357, P158
- (26) Martemyanov, K; FEBS Lett 1997, V414, P268 HCAPLUS
- (27) Martemyanov, K; Protein Expression and Purification, in press 2001
- (28) Nirenberg, M; Proc Natl Acad Sci USA 1961, V47, P1588 HCAPLUS
- (29) Nishimura, N; Biotech Lett 1993, V15, P785 HCAPLUS
- (30) Nishimura, N; J Ferment Bioeng 1995, V79, P131 HCAPLUS
- (31) Ryabova, L; J Biol Chem 1994, V269, P1501 HCAPLUS
- (32) Ryabova, L; Methods in Molecular Biology, Protein Synthesis: Methods and Protocols 1998, V77, P179 HCAPLUS
- (33) Ryabova, L; Nature Biotechnology 1997, V15, P79 HCAPLUS
- (34) Ryabova, L; Nucleic Acids Res 1989, V17, P4412 HCAPLUS
- (35) Spirin, A; Frontiers in Bioprocessing 1991, VII, P31
- (36) Spirin, A; Ribosomes 1999
- (37) Spirin, A; Science 1988, V242, P1162 HCAPLUS
- (38) Uzawa, T; J Biochem (Japan) 1993, V114, P732 HCAPLUS
- (39) Volyanik, E; Anal Biochem 1993, V214, P289 HCAPLUS

L60 ANSWER 10 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:29500 HCAPLUS

DN 138:84441

TI PCR amplification of template DNA for **cell-free protein synthesis** system

IN Motoda, Yoko; Yabuki, Takashi; Kikawa, Takanori; Yokoyama, Shigeyuki

PA Institute of Physical and Chemical Research, Japan

SO Jpn. Kokai Tokkyo Koho, 17 pp.

CODEN: JKXXAF

DT **Patent**

LA Japanese

IC ICM C12N015-09

ICS C12P021-02

CC 3-1 (Biochemical Genetics)

Section cross-reference(s): 6, 9

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2003009880	A2	20030114	JP 2001-201356	20010702
	WO 2003004703	A1	20030116	WO 2002-JP6261	20020624
	W: CA, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				

PRAI JP 2001-201356 A 20010702

AB A process for producing template DNA for use in **cell-free protein synthesis** by DNA amplification, is provided. A first double-stranded DNA (dsDNA) coding for a **protein**, a second dsDNA overlapping at the 5' end, a third dsDNA overlapping at the 3' end, a sense primer hybridizable to the 5' end of the second dsDNA, an antisense primer hybridizable to the 3' end of the 3rd dsDNA, are used for polymerase chain reaction (PCR). Single-stranded DNA can also be used in place of dsDNA. Use of a terminator sequence and peptide tag such as maltose-binding **protein**, cellulose-binding **domain**, glutathione-S-transferase, thioredoxin, streptavidin-binding peptide, or histidine-contg. peptide, is claimed. **Expression of Ras protein in cell-free protein synthesis** system using the native histidine tag present in the N-terminal of chicken muscle lactate dehydrogenase, T7 promoter, and T7 terminator, is described. Chloramphenicol acetyltransferase, and 10 arbitrarily chosen **clones** from mouse full-length cDNA library were also **expressed**. By using this process, a desired polypeptide can be **synthesized** within a short period of time at a high yield and at a low cost, compared with the

conventional processes.

- ST PCR amplification template DNA cell translation system
- IT Nucleic acid amplification (method)  
(DNA; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT PCR (polymerase chain reaction)  
**Protein** sequences  
(PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT **Proteins**  
Ras **proteins**  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT Primers (nucleic acid)  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT **Translation, genetic**  
(**cell-free**; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT DNA  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(double-stranded; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT Peptides, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(histidine-contg., use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT DNA  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(single-stranded; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT Peptides, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(streptavidin-binding, use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT Peptides, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(tag; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT Genetic element  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(terminator; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT Thioredoxins  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT 9040-07-7P, Chloramphenicol acetyltransferase  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP

(Preparation)

(PCR amplification of template DNA for **cell-free protein synthesis** system)

IT 140295-47-2, GenBank M21532 140531-55-1, GenBank M16206 142318-93-2,  
GenBank X65627 150947-97-0, GenBank L16904 175301-96-9, GenBank U51204  
180770-00-7, GenBank D87663 186016-45-5, GenBank U85511 252790-02-6,  
GenBank S68022 384441-00-3, GenBank X13605 391530-17-9, GenBank M32599  
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL  
(Biological study)

(PCR amplification of template DNA for **cell-free protein synthesis** system)

IT 484053-89-6  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)

(amino acid sequence; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT 484100-59-6 484100-60-9 484100-61-0 484100-62-1 484100-63-2  
484100-64-3 484100-65-4 484100-66-5 484100-67-6 484100-68-7  
484100-69-8 484100-70-1 484100-71-2 484100-72-3 484100-73-4  
484100-74-5 484100-75-6 484100-76-7 484100-77-8 484100-78-9  
484100-79-0 484100-80-3 484100-81-4

RL: PRP (Properties)

(unclaimed nucleotide sequence; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT 50812-37-8, Glutathione S-transferase  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)

(use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)

L60 ANSWER 11 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2003:29497 HCAPLUS

DN 138:84439

TI PCR amplification of template DNA for **cell-free protein synthesis** system

IN Motoda, Yoko; Yabuki, Takashi; Kikawa, Takanori; Yokoyama, Shigeyuki

PA Institute of Physical and Chemical Research, Japan

SO Jpn. Kokai Tokkyo Koho, 17 pp.

CODEN: JKXXAF

DT **Patent**

LA Japanese

IC ICM C12N015-09

ICS C12P021-02

CC 3-1 (Biochemical Genetics)

Section cross-reference(s): 6, 9

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2003009877	A2	20030114	JP 2001-200676	20010702
PRAI	JP 2001-200676		20010702		

AB A process for producing template DNA for use in **cell-free protein synthesis** by DNA amplification, is provided. A first double-stranded DNA (dsDNA) coding for a **protein**, a second dsDNA overlapping at the 5' end, a third dsDNA overlapping at the 3' end, a sense primer hybridizable to the 5' end of the second dsDNA, an antisense primer hybridizable to the 3' end of the 3rd dsDNA, are used for polymerase chain reaction (PCR). Single-stranded DNA can also be used in place of dsDNA. Use of a terminator sequence and peptide tag such as maltose-binding **protein**, cellulose-binding **domain**, glutathione-S-transferase, thioredoxin,

streptavidin-binding peptide, or histidine-contg. peptide, is claimed.

**Expression of Ras protein in cell-free**

**protein synthesis** system using the native histidine tag present in the N-terminal of chicken muscle lactate dehydrogenase, T7 promoter, and T7 terminator, is described. Chloramphenicol acetyltransferase, and 10 arbitrarily chosen **clones** from mouse full-length cDNA library were also **expressed**. By using this process, a desired polypeptide can be **synthesized** within a short period of time at a high yield and at a low cost, compared with the conventional processes.

ST PCR amplification template DNA cell translation system

IT Nucleic acid amplification (method)

(DNA; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT PCR (polymerase chain reaction)

**Protein** sequences

(PCR amplification of template DNA for **cell-free protein synthesis** system)

IT **Proteins**

Ras **proteins**

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

(PCR amplification of template DNA for **cell-free protein synthesis** system)

IT Primers (nucleic acid)

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(PCR amplification of template DNA for **cell-free protein synthesis** system)

IT **Translation, genetic**

(**cell-free**; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT DNA

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(double-stranded; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT Peptides, biological studies

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(histidine-contg., use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT DNA

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(single-stranded; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT Peptides, biological studies

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(streptavidin-binding, use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT Peptides, biological studies

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(tag; PCR amplification of template DNA for **cell-free protein synthesis** system)

IT Genetic element

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

- (terminator; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT Thioredoxins  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT 9040-07-7P, Chloramphenicol acetyltransferase  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP  
(Preparation)  
(PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT 140295-47-2, GenBank M21532 140531-55-1, GenBank M16206 142318-93-2, GenBank X65627 150947-97-0, GenBank L16904 175301-96-9, GenBank U51204 180770-00-7, GenBank D87663 186016-45-5, GenBank U85511 252790-02-6, GenBank S68022 384441-00-3, GenBank X13605 391530-17-9, GenBank M32599  
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL  
(Biological study)  
(PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT 484053-89-6  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(amino acid sequence; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT 484100-82-5 484100-83-6 484100-84-7 484100-85-8 484100-86-9  
484100-87-0 484100-88-1 484100-89-2 484100-90-5 484100-91-6  
484100-92-7 484100-93-8 484100-94-9 484100-95-0 484100-96-1  
484100-97-2 484100-98-3 484100-99-4 484101-00-0 484101-01-1  
484101-02-2 484101-03-3 484101-04-4  
RL: PRP (Properties)  
(unclaimed nucleotide sequence; PCR amplification of template DNA for **cell-free protein synthesis** system)
- IT 50812-37-8, Glutathione S-transferase  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)  
(use as tag; PCR amplification of template DNA for **cell-free protein synthesis** system)
- L60 ANSWER 12 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2003:11293 HCAPLUS  
DN 138:299991  
TI Selenomethionine incorporation into a **protein** by **cell-free synthesis**
- AU **Kigawa, Takanori**; Yamaguchi-Nunokawa, Emi; Kodama, Koichiro; Matsuda, Takayoshi; Yabuki, Takashi; Matsuda, Natsuko; Ishitani, Ryuichiro; Nureki, Osamu; **Yokoyama, Shigeyuki**
- CS RIKEN Genomic Sciences Center, Yokohama, 230-0045, Japan  
SO Journal of Structural and Functional Genomics (2002), 2(1), 29-35  
CODEN: JSFGAW; ISSN: 1345-711X  
PB Kluwer Academic Publishers  
DT Journal  
LA English  
CC **9-5 (Biochemical Methods)**  
Section cross-reference(s): 6
- AB Multi-wavelength anomalous diffraction phasing is esp. useful for high-throughput structure detns. Selenomethionine substituted **proteins** are commonly used for this purpose. However, the cytotoxicity of selenomethionine drastically reduces the efficiency of its incorporation in in vivo **expression** systems. In the present study, an improved **E. coli cell-free**

**protein synthesis** system was used to incorporate selenomethionine into a **protein**, so that highly efficient incorporation could be achieved. A milligram quantity of selenomethionine-contg. Ras was obtained using the **cell-free** system with dialysis. The mass spectrometry anal. showed that more than 95% of the methionine residues were substituted with selenomethionine. The crystal of this **protein** grew under the same conditions and had the same unit **cell** constns. as those of the native Ras **protein**. The three-dimensional structure of this **protein**, detd. by multi-wavelength anomalous diffraction phasing, was almost the same as that of the Ras **protein** prepd. by in vivo **expression**. Therefore, the **cell-free synthesis** system could become a powerful **protein expression** method for high-throughput structure detns. by x-ray crystallog.

ST selenomethionine **protein** cell **synthesis**

IT Laser ionization mass spectrometry  
(photodesorption, matrix-assisted; selenomethionine incorporation into **protein** by **cell-free synthesis**)

IT Laser desorption mass spectrometry  
(photoionization, matrix-assisted; selenomethionine incorporation into **protein** by **cell-free synthesis**)

IT **Conformation**  
(**protein**; selenomethionine incorporation into **protein** by **cell-free synthesis**)

IT Ras **proteins**  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(selenomethionine contg.; selenomethionine incorporation into **protein** by **cell-free synthesis**)

IT **Escherichia coli**  
Time-of-flight mass spectrometry  
(selenomethionine incorporation into **protein** by **cell-free synthesis**)

IT **Proteins**  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(selenomethionine incorporation into **protein** by **cell-free synthesis**)

IT Crystallography  
(x-ray; selenomethionine incorporation into **protein** by **cell-free synthesis**)

IT 1464-42-2, Selenomethionine  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(selenomethionine incorporation into **protein** by **cell-free synthesis**)

RE.CNT 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Akashi, S; J Mass Spectrom Soc Jpn 1996, V44, P269 HCAPLUS
- (2) Brunger, A; Acta Crystallogr 1998, VD54, P905 HCAPLUS
- (3) Cayanis, E; J Biol Chem 1990, V265, P10829 HCAPLUS
- (4) Collaborative Computational Project Number 4; Acta Crystallogr 1994, VD50, P760
- (5) Davis, J; Promega Notes Magazine 1996, V56, P14
- (6) de La Fortelle, E; Methods Enzymol 1997, V276, P472 HCAPLUS
- (7) de Vos, A; Science 1988, V239, P888 HCAPLUS
- (8) Ha, J; Biochemistry 1989, V28, P8411 HCAPLUS
- (9) Hauptman, H; Acta Crystallogr 1999, V55, P891
- (10) Hendrickson, W; EMBO J 1990, V9, P1665 HCAPLUS
- (11) Hendrickson, W; Science 1991, V254, P51 HCAPLUS
- (12) Ito, Y; Biochemistry 1997, V36, P9109 HCAPLUS
- (13) Jones, T; Acta Crystallogr 1991, VA47, P110 HCAPLUS
- (14) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (15) Kigawa, T; J Biomol NMR 1995, V6, P129 HCAPLUS
- (16) Kim, D; Biotechnol Prog 1996, V12, P645 HCAPLUS



- (17) Kim, D; Eur J Biochem 1996, V239, P881 HCAPLUS
- (18) Knight, S; Acta Crystallogr 2000, VD56, P42 HCAPLUS
- (19) Koradi, R; J Mol Graph 1996, V14, P51 HCAPLUS
- (20) Kraulis, P; Biochemistry 1994, V33, P3515 HCAPLUS
- (21) Madin, K; Proc Natl Acad Sci U S A 2000, V97, P559 HCAPLUS
- (22) Markoff, L; J Virol 1989, V63, P3345 HCAPLUS
- (23) Martemyanov, K; Protein Expr Purif 2001, V21, P456 HCAPLUS
- (24) Otwinowski, Z; Macromol Crystallogr 1997, VA276, P307
- (25) O'Gara, M; Eur J Biochem 1997, V247, P1009 HCAPLUS
- (26) Pratt, J; Transcription and translation 1984, P179 HCAPLUS
- (27) Ryabova, L; Nat Biotechnol 1997, V15, P79 HCAPLUS
- (28) Tsalkova, T; Biochemistry 1993, V32, P3377 HCAPLUS
- (29) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS
- (30) Yamasaki, K; Biochem Biophys Res Commun 1989, V162, P1054 HCAPLUS
- (31) Yokoyama, S; Nat Struct Biol 2000, V7(Suppl), P943
- (32) Zawadzki, V; Nucleic Acids Res 1991, V19, P1948 HCAPLUS

L60 ANSWER 13 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:946589 HCAPLUS

DN 138:21811

TI Method for analyzing **protein** structure by mass spectrometry

IN Shen, Xinchun; Kigawa, Takanori; Yokoyama, Shigeyuki

PA Riken Corp., Japan

SO PCT Int. Appl., 35 pp.

CODEN: PIXXD2

DT **Patent**

LA Japanese

IC ICM G01N033-68

ICS G01N027-62; G01N033-53

CC **9-16 (Biochemical Methods)**

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	WO 2002099437	A1	20021212	WO 2002-JP4473	20020508
	W: CA, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
	JP 2002372517	A2	20021226	JP 2001-164819	20010531
PRAI	JP 2001-164819	A	20010531		
AB	A method is provided for analyzing <b>protein</b> mol. wt. by mass spectrometry. The method comprises: (a) a step for <b>synthesizing</b> a <b>protein fused</b> with a tag sequence with a <b>cell-free protein synthesis</b> system; (b) a step for bringing the <b>fused protein synthesized</b> into contact with a carrier capable of specifically binding to the tag sequence; (c) a step for recovering of the carrier bound to the <b>fused protein</b> ; and (d) a step for measuring the mol. wt. of the <b>fused protein</b> by mass spectrometry (e.g., MALDI-TOF) using the <b>fused protein</b> bound to the carrier recovered. By using this method, a series of procedures of <b>synthesizing</b> , purifying and analyzing a <b>protein</b> can be efficiently and quickly carried out.				
ST	<b>protein</b> tag sequence carrier mass spectrometry				
IT	<b>Translation, genetic</b> ( <b>cell-free</b> ; method for analyzing <b>protein</b> structure by mass spectroscopy)				
IT	<b>Proteins</b> RL: NUU (Other use, unclassified); USES (Uses) (ligand-binding; method for analyzing <b>protein</b> structure by mass spectroscopy)				
IT	Carriers Mass spectrometry Molecular weight				

**Protein motifs**(method for analyzing **protein** structure by mass spectroscopy)IT **Proteins**

RL: ANT (Analyte); BSU (Biological study, unclassified); PRP (Properties); ANST (Analytical study); BIOL (Biological study)

(method for analyzing **protein** structure by mass spectroscopy)

## IT Amino acids, uses

Antibodies

Heterocyclic compounds

Metals, uses

Oligonucleotides

Peptides, uses

Receptors

RL: NUU (Other use, unclassified); USES (Uses)

(method for analyzing **protein** structure by mass spectroscopy)

## IT Laser ionization mass spectrometry

(photodesorption, matrix-assisted, TOF; method for analyzing **protein** structure by mass spectroscopy)

## IT Laser desorption mass spectrometry

(photoionization, matrix-assisted, TOF; method for analyzing **protein** structure by mass spectroscopy)

## IT 67663-36-9, Ras

RL: ANT (Analyte); BSU (Biological study, unclassified); PRP (Properties); ANST (Analytical study); BIOL (Biological study)

(fused with a histidine tag; method for analyzing **protein** structure by mass spectroscopy)

## IT 9001-92-7, Proteinase

RL: NUU (Other use, unclassified); USES (Uses)

(method for analyzing **protein** structure by mass spectroscopy)

## IT 9012-36-6, Agarose

RL: NUU (Other use, unclassified); USES (Uses)

(nickel-bound beads; method for analyzing **protein** structure by mass spectroscopy)

## IT 478220-36-9

RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)

(unclaimed amino acid sequence)

## IT 478220-37-0

RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)

(unclaimed nucleotide sequence)

IT 478235-80-2 478235-82-4 478235-84-6 478235-85-7, 8: PN: WO02099437  
SEQID: 4 unclaimed DNA 478237-90-0, 9: PN: WO02099437 SEQID: 5 unclaimed DNA

RL: PRP (Properties)

(unclaimed nucleotide sequence; method for analyzing **protein** structure by mass spectrometry)

## IT 478235-81-3 478235-83-5

RL: PRP (Properties)

(unclaimed **protein** sequence; method for analyzing **protein** structure by mass spectrometry)RE.CNT 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Max Planck Ges Foerderung Wissenschaften; EP 1036202 A 1999 HCAPLUS
- (2) Max Planck Ges Foerderung Wissenschaften; JP 2001526381 A 1999
- (3) Max Planck Ges Foerderung Wissenschaften; WO 9929898 A 1999 HCAPLUS
- (4) Pharmacia Biosensor Ab; JP 11-512518 A 1999
- (5) Pharmacia Biosensor Ab; US 5955729 A 1999 HCAPLUS
- (6) Pharmacia Biosensor Ab; EP 850407 A 1999 HCAPLUS
- (7) Pharmacia Biosensor Ab; WO 9709608 A 1999 HCAPLUS
- (8) Rikagaku, K; WO 0036133 A 2000 HCAPLUS
- (9) Rikagaku, K; EP 1143009 A 2000 HCAPLUS
- (10) Rikagaku, K; JP 2000175695 A 2000 HCAPLUS

- (11) Rikagaku, K; WO 0068690 A 2001 HCAPLUS
- (12) Rikagaku, K; EP 1182458 A 2001 HCAPLUS
- (13) Rikagaku, K; JP 200127633 A 2001
- (14) Sequenom Inc; EP 1010008 A 1999 HCAPLUS
- (15) Sequenom Inc; DE 19882657 T 1999 HCAPLUS
- (16) Sequenom Inc; JP 2001515209 A 1999
- (17) Sequenom Inc; US 6207370 B1 1999 HCAPLUS
- (18) Sequenom Inc; WO 9912040 A 1999 HCAPLUS

L60 ANSWER 14 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:869077 HCAPLUS

DN 137:365990

TI Detergents used for biosynthesis of membrane **protein** in  
**cell-free protein synthesis** system

IN Tajima, Kaori; Kigawa, Takanori; Shirouzu, Mikako; Yabuki,  
Takashi; Ishihara, Goushi; Yokoyama, Shigeyuki

PA Riken Corp., Japan

SO PCT Int. Appl., 45 pp.

CODEN: PIXXD2

DT **Patent**

LA Japanese

IC ICM C12N015-09

ICS C12P021-02; C12N001-21; C07K001-02

CC **9-14 (Biochemical Methods)**

Section cross-reference(s): 3, 6

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002090537	A1	20021114	WO 2002-JP4204	20020426
	W: CA, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
	JP 2003018999	A2	20030121	JP 2001-285145	20010919
PRAI	JP 2001-135111	A	20010502		
	JP 2001-285145	A	20010919		
AB	In a process for producing a <b>protein</b> using a <b>cell-free protein synthesis</b> system, the system contains a detergent so that the <b>protein</b> can be <b>synthesized</b> without aggregation and denaturation. The above-described <b>protein</b> is a <b>protein</b> having a hydrophobic <b>domain</b> at least a part thereof, for example, a membrane <b>protein</b> or its fragment. The concn. of digitonin and Brij35 was 1-2 % and 0.01-0.5 %, resp. in the reaction buffer. The above-described detergent is a mild detergent which would not denature the <b>protein</b> , for example, a nonionic or amphoteric detergent.				
ST	detergent membrane <b>protein synthesis cell free</b> system				
IT	Polyoxyalkylenes, biological studies				
	RL: BSU (Biological study, unclassified); BIOL (Biological study) (alkyl ethers; detergents used for biosynthesis of membrane <b>protein in cell-free protein synthesis</b> system)				
IT	Detergents (ampholytic; detergents used for biosynthesis of membrane <b>protein in cell-free protein synthesis</b> system)				
IT	<b>Proteins</b>				
	RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation) (channel, <b>synthesis</b> of; detergents used for biosynthesis of membrane <b>protein in cell-free protein synthesis</b> system)				
IT	<b>Proteins</b>				

- RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(hydrophobic, **synthesis** of; detergents used for biosynthesis  
of membrane **protein** in **cell-free**  
**protein synthesis** system)
- IT DNA  
Lipids, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(in reaction buffer; detergents used for biosynthesis of membrane  
**protein** in **cell-free protein**  
**synthesis** system)
- IT **Proteins**  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(membrane, hydrophobic, **synthesis** of; detergents used for  
biosynthesis of membrane **protein** in **cell-**  
**free protein synthesis** system)
- IT Detergents  
(nonionic; detergents used for biosynthesis of membrane **protein**  
in **cell-free protein synthesis**  
system)
- IT **Translation, genetic**  
(of membrane **protein**, method of; detergents used for  
biosynthesis of membrane **protein** in **cell-**  
**free protein synthesis** system)
- IT Aggregation  
(of **protein**, prevention of; detergents used for biosynthesis  
of membrane **protein** in **cell-free**  
**protein synthesis** system)
- IT Denaturation  
(prevention of; detergents used for biosynthesis of membrane  
**protein** in **cell-free protein**  
**synthesis** system)
- IT Receptors  
Transport **proteins**  
RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified);  
BIOL (Biological study); PREP (Preparation)  
(**synthesis** of; detergents used for biosynthesis of membrane  
**protein** in **cell-free protein**  
**synthesis** system)
- IT 2605-79-0, N,N-Dimethyldecylamine N-oxide 7305-30-8 9002-92-0, Brij35  
9005-63-4, Polyoxyethylene sorbitan 11024-24-1, Digitonin 15163-36-7  
19327-40-3, 3,6,9,12,15-Pentaoxatricosan-1-ol 25322-68-3D, alkyl ethers  
29836-26-8, Octyl glucoside 31835-06-0, Sucrose mono decanoate  
69227-93-6, .beta.-Dodecyl maltoside 75621-03-3, Chaps 85261-19-4,  
Nonanoyl-N-methyl glucamide 85261-20-7, Decanoyl-N-methyl glucamide  
85618-20-8, .beta.-Heptylthioglucoiside 85618-21-9, .beta.-  
Octylthioglucoiside  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(detergents used for biosynthesis of membrane **protein** in  
**cell-free protein synthesis**  
system)
- IT 475460-14-1, 1: PN: WO02090537 SEQID: 1 unclaimed DNA 475460-15-2  
475460-16-3, 3: PN: WO02090537 SEQID: 3 unclaimed DNA 475460-17-4, 4:  
PN: WO02090537 SEQID: 4 unclaimed DNA 475460-18-5 475460-19-6  
475460-20-9, 7: PN: WO02090537 SEQID: 7 unclaimed DNA 475460-21-0, 8:  
PN: WO02090537 SEQID: 8 unclaimed DNA  
RL: PRP (Properties)  
(unclaimed nucleotide sequence; detergents used for biosynthesis of  
membrane **protein** in **cell-free**  
**protein synthesis** system)

RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Booth, P; Biochemical Society Transactions, part 3 2000, V28, PA50  
(2) Kigawa, T; FEBS Letters 1999, V442, P15 HCAPLUS

- (3) Medical Research Council; EP 1019496 A1 1999 HCAPLUS
- (4) Medical Research Council; JP 2001509380 A 1999
- (5) Medical Research Council; WO 9902671 A1 1999 HCAPLUS
- (6) Nishimura, N; Journal of Fermentation and Bioengineering 1995, V80(4), P403 HCAPLUS
- (7) Rikagaku Kabushiki Kaisha; WO 0036133 A1 2000 HCAPLUS
- (8) Rikagaku Kabushiki Kaisha; EP 1143009 A1 2000 HCAPLUS
- (9) Rikagaku Kabushiki Kaisha; JP 2000175695 A 2000 HCAPLUS
- (10) Wakenyaku Co Ltd; WO 0068412 A1 2000 HCAPLUS
- (11) Wakenyaku Co Ltd; EP 1176210 A1 2000 HCAPLUS
- (12) Wakenyaku Co Ltd; JP 2000316594 A 2000 HCAPLUS

L60 ANSWER 15 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:704583 HCAPLUS

DN 137:227659

TI **Cell-free synthesis of soluble protein domains for three-dimensional structure analysis**

IN **Seki, Eiko; Kikawa, Takanori; Yokoyama, Shigeyuki**

PA Institute of Physical and Chemical Research, Japan

SO Jpn. Kokai Tokkyo Koho, 9 pp.

CODEN: JKXXAF

DT **Patent**

LA Japanese

IC ICM C12N015-09

ICS C07K014-00; C07K019-00; C12N001-21; C12P021-02; G01N023-20; G01N024-08; G01N033-68; C12R001-19

CC **3-2 (Biochemical Genetics)**

Section cross-reference(s): 9

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2002262873	A2	20020917	JP 2001-62703	20010306
	US 2002142387	A1	20021003	US 2001-994573	20011126
PRAI	JP 2001-62703	A	20010306		

AB A method for **cell-free synthesis** of **sol. protein domains** suitable for **three-dimensional structure anal.**, is disclosed. **Protein domains** are **expressed** as **fusion proteins** with a detectable **protein** such as **enzyme**, binding **protein**, luminescent **protein**, or fluorescent **protein**, and selected. They can be used for structure detn. by x-ray crystallog. or NMR. **Expression** of mammalian growth hormone receptor binding **protein Grb2 domains** as **fusion protein** with **GFP** mutant is described. Fragments having **domain** structure at the N-terminal produced stronger fluorescence than those having **domain** structure in the middle.

ST **cell free synthesis protein**

IT **domain three dimensional structure analysis**

IT **Proteins**

RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified);

BIOL (Biological study); PREP (Preparation)

(Grb-2, **domain expression** of; **cell-**

**free synthesis** of **sol. protein**

**domains** for **three-dimensional structure anal.**)

IT **Molecular structure determination methods**

(NMR spectrometric; **cell-free synthesis**

of **sol. protein domains** for **three**

**-dimensional structure anal.**)

IT **Crystal structure determination methods**

-- Applicant

**Protein motifs**

(cell-free synthesis of sol.  
protein domains for three-  
dimensional structure anal.)

IT **Fusion proteins (chimeric proteins)**

RL: ARU (Analytical role, unclassified); BPN (Biosynthetic preparation);  
BUU (Biological use, unclassified); ANST (Analytical study); BIOL  
(Biological study); PREP (Preparation); USES (Uses)  
(cell-free synthesis of sol.  
protein domains for three-  
dimensional structure anal.)

IT **Translation, genetic**

(cell-free; cell-free  
synthesis of sol. protein domains  
for three-dimensional structure anal.)

IT **Proteins**

RL: ARU (Analytical role, unclassified); BPN (Biosynthetic preparation);  
BUU (Biological use, unclassified); ANST (Analytical study); BIOL  
(Biological study); PREP (Preparation); USES (Uses)  
(green fluorescent, fusion  
protein of; cell-free synthesis  
of sol. protein domains for three  
-dimensional structure anal.)

IT **Conformation**

**Quaternary structure**

**Tertiary structure**

(protein; cell-free synthesis  
of sol. protein domains for three  
-dimensional structure anal.)

IT **Molecular structure determination methods**

(x-ray diffractometric; cell-free synthesis  
of sol. protein domains for three  
-dimensional structure anal.)

## IT 459469-42-2

RL: PRP (Properties)  
(unclaimed nucleotide sequence; cell-free  
synthesis of sol. protein domains  
for three-dimensional structure anal.)

L60 ANSWER 16 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:704578 HCAPLUS

DN 137:212639

TI **Cell-free synthesis** of heavy atom-containing  
**proteins** for x-ray crystallography structural analysis

IN Nunokawa, Emi; Kikawa, Takanori; Yabuki, Takashi; Yokoyama,  
Shigeyuki

PA Institute of Physical and Chemical Research, Japan

SO Jpn. Kokai Tokkyo Koho, 10 pp.

CODEN: JKXXAF

DT **Patent**

LA Japanese

IC ICM C12N015-09

ICS C07K001-34; C12P021-00; C12N001-16; C12N001-20

CC 6-3 (General Biochemistry)

Section cross-reference(s): 9, 75

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	JP 2002262867	A2	20020917	JP 2001-65799	20010308
	US 2002168705	A1	20021114	US 2001-989974	20011120
PRAI	JP 2001-65799	A	20010308		
AB	A method for large-scale <b>cell-free synthesis</b> of heavy atom-contg. <b>proteins</b> suitable for x-ray crystallog.				

structural anal. using dialysis, is disclosed. **Cell** ext. of **E. coli**, hyperthermophilic archaeon, or yeast, is used. It also includes ATP regeneration system, macromol. adsorbent, and reducing agent. Creatine kinase and creatine phosphate are used for ATP regeneration. Amino acids contg. mercury, platinum, iodine, iron, or selenium, such as selenocysteine or selenomethionine, are to be incorporated. **Synthesis** of selenomethionine-contg. **Ras protein** by **cell-free synthesis** system, crystn. by hanging-drop vapor-diffusion method, and structural anal. by multiwavelength anomalous diffraction (MAD), are described. The three dimensional structure model produced was identical to those of unlabeled **proteins** produced in vivo and in **cell-free** system.

- ST **cell free protein synthesis** heavy atom replacement xray crystallog
- IT Archaeobacteria (Archaea)
  - Escherichia coli**
  - Yeast
    - (cell ext. of; **cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT **Crystal structure determination methods**
  - Dialysis
  - Reducing agents
    - (**cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT **Ras proteins**
  - RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)
  - (**cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT **Translation, genetic**
  - (**cell-free; cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT Amino acids, biological studies
  - RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
  - (heavy atom-contg.; **cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT **Conformation**
  - Quaternary structure**
  - Tertiary structure**
    - (**protein; cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT **Molecular structure determination methods**
  - (x-ray diffractometric; **cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT 7439-89-6, Iron, biological studies 7439-97-6, Mercury, biological studies 7440-06-4, Platinum, biological studies 7553-56-2, Iodine, biological studies 7782-49-2, Selenium, biological studies
  - RL: ARU (Analytical role, unclassified); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
  - (amino acids contg.; **cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT 3211-76-5, Selenomethionine 3614-08-2, Selenocysteine
  - RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

- (**cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT 56-65-5, 5'-ATP, biological studies  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(regeneration system for; **cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT 67-07-2, Creatine phosphate  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(use for ATP regeneration; **cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- IT 9001-15-4, Creatine kinase  
RL: BUU (Biological use, unclassified); CAT (Catalyst use); BIOL (Biological study); USES (Uses)  
(use for ATP regeneration; **cell-free synthesis** of heavy atom-contg. **proteins** for x-ray crystallog. structural anal.)
- L60 ANSWER 17 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2002:576300 HCAPLUS  
DN 137:258909  
TI An engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of an unnatural amino acid into **proteins** in eukaryotic translation and its application in a wheat germ **cell-free** system
- AU Kiga, Daisuke; Sakamoto, Kensaku; Kodama, Koichiro; **Kigawa, Takanori**; Matsuda, Takayoshi; Yabuki, Takashi; Shirouzu, Mikako; Harada, Yoko; Nakayama, Hiroshi; Takio, Koji; Hasegawa, Yoshinori; Endo, Yaeta; Hirao, Ichiro; **Yokoyama, Shigeyuki**
- CS RIKEN Genomic Sciences Center, Yokohama, 230-0045, Japan  
SO Proceedings of the National Academy of Sciences of the United States of America (2002), 99(15), 9715-9723  
CODEN: PNASA6; ISSN: 0027-8424
- PB National Academy of Sciences  
DT Journal  
LA English  
CC 6-1 (General Biochemistry)  
Section cross-reference(s): 7
- AB Tyrosyl-tRNA **synthetase** (TyrRS) from *Escherichia coli* was engineered to preferentially recognize 3-iodo-L-tyrosine rather than L-tyrosine for the site-specific incorporation of 3-iodo-L-tyrosine into **proteins** in eukaryotic translation systems. The wild-type TyrRS does not recognize 3-iodo-L-tyrosine, because of the bulky iodine substitution. On the basis of the reported crystal structure of *Bacillus stearothermophilus* TyrRS, three residues, Y37, Q179, and Q195, in the L-tyrosine-binding site were chosen for mutagenesis. Thirty-four single amino acid replacements and 16 of their combinations were screened by in vitro biochem. assays. A combination of the Y37V and Q195C mutations changed the amino acid specificity in such a way that the variant TyrRS activates 3-iodo-L-tyrosine 10-fold more efficiently than L-tyrosine. This engineered **enzyme**, TyrRS(V37C195), was tested for use in the wheat germ **cell-free** translation system, which has recently been significantly improved, and is now as productive as conventional **recombinant** systems. During the translation in the wheat germ system, an *E. coli* suppressor tRNA<sup>Tyr</sup> was not aminoacylated by the wheat germ **enzymes**, but was aminoacylated by the *E. coli* TyrRS(V37C195) variant with 3-iodo-L-tyrosine. After the use of the 3-iodotyrosyl-tRNA in translation, the resultant uncharged tRNA could be aminoacylated again in the system. A mass spectrometric anal. of the



produced **protein** revealed that more than 95% of the amino acids incorporated for an amber codon were iodotyrosine, whose concn. was only twice that of L-tyrosine in the translation. Therefore, the variant **enzyme**, 3-iodo-L-tyrosine, and the suppressor tRNA can serve as an addnl. set orthogonal to the 20 endogenous sets in eukaryotic in vitro translation systems.

ST engineering *Escherichia* tyrosyl tRNA **synthetase** unnatural amino acid translation

IT Codons

RL: BSU (Biological study, unclassified); BIOL (Biological study) (amber; engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT **Enzyme kinetics**

**Protein engineering**

**Translation, genetic**

(engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT **Enzyme functional sites**

(substrate-binding, tyrosine-binding; engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT tRNA

RL: BSU (Biological study, unclassified); BIOL (Biological study) (suppressor; engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT 9023-45-4P, Tyrosyl-tRNA **synthetase**

RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation) (engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT 60-18-4, L-Tyrosine, biological studies 70-78-0, 3-Iodo-L-tyrosine

RL: BSU (Biological study, unclassified); BIOL (Biological study) (engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT 52-90-4, L-Cysteine, biological studies

RL: BSU (Biological study, unclassified); BIOL (Biological study) (residue no. 195 in mutant **synthetase**; engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT 56-85-9, L-Glutamine, biological studies

RL: BSU (Biological study, unclassified); BIOL (Biological study) (residue no. 195 in wild-type **synthetase**; engineered *Escherichia coli* tyrosyl-tRNA **synthetase** for site-specific incorporation of unnatural amino acid into **proteins** in eukaryotic translation and application in a wheat germ **cell-free** system)

IT 72-18-4, L-Valine, biological studies

RL: BSU (Biological study, unclassified); BIOL (Biological study) (residue no. 37 in mutant **synthetase**; engineered

**Escherichia coli** tyrosyl-tRNA synthetase  
for site-specific incorporation of unnatural amino acid into  
**proteins** in eukaryotic translation and application in a wheat  
germ **cell-free** system)

RE.CNT 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Bain, J; J Am Chem Soc 1989, V111, P8013 HCAPLUS
- (2) Bain, J; Nature 1992, V356, P537 HCAPLUS
- (3) Brick, P; J Mol Biol 1989, V208, P83 HCAPLUS
- (4) Calendar, R; Biochemistry 1966, V5, P1690 HCAPLUS
- (5) Doering, V; Science 2001, V292, P501
- (6) Drabkin, H; Mol Cell Biol 1996, V16, P907 HCAPLUS
- (7) Edwards, H; Mol Cell Biol 1990, V10, P1633 HCAPLUS
- (8) Edwards, H; Proc Natl Acad Sci 1991, V88, P1153 HCAPLUS
- (9) Furter, R; Prot Sci 1998, V7, P419 HCAPLUS
- (10) Ha, J; Biochemistry 1989, V28, P8411 HCAPLUS
- (11) Hamano-Takaku, F; J Biol Chem 2000, V275, P40324 HCAPLUS
- (12) Hecht, S; Acc Chem Res 1992, V25, P545 HCAPLUS
- (13) Hecht, S; J Biol Chem 1978, V253, P4517 HCAPLUS
- (14) Hendrickson, W; EMBO J 1990, V9, P1665 HCAPLUS
- (15) Hirao, I; Nat Biotechnol 2002, V20, P177 HCAPLUS
- (16) Kamada, K; Nature 1996, V383, P598 HCAPLUS
- (17) Kast, P; J Mol Biol 1991, V222, P99 HCAPLUS
- (18) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (19) Kigawa, T; J Struct Funct Genomics 2001, V2, P27
- (20) Kiick, K; FEBS Lett 2001, V502, P25 HCAPLUS
- (21) Kim, D; Biotechnol Bioeng 2001, V74, P309 HCAPLUS
- (22) Kohrer, C; Proc Natl Acad Sci 2001, V98, P14310 HCAPLUS
- (23) Koide, H; Biochemistry 1994, V33, P7470 HCAPLUS
- (24) Koide, H; Proc Natl Acad Sci 1988, V85, P6237 HCAPLUS
- (25) Kowal, A; Proc Natl Acad Sci 2001, V98, P2268 HCAPLUS
- (26) Lloyd, A; Nucleic Acids Res 1995, V23, P2886 HCAPLUS
- (27) Lowe, D; Biochemistry 1987, V26, P6038 HCAPLUS
- (28) Madin, K; Proc Natl Acad Sci 2000, V97, P559 HCAPLUS
- (29) Moore, B; J Mol Biol 2000, V298, P195 HCAPLUS
- (30) Murakami, H; J Am Chem Soc 1998, V120, P7520 HCAPLUS
- (31) Noren, C; Science 1989, V244, P182 HCAPLUS
- (32) Nowak, M; Science 1995, V268, P439 HCAPLUS
- (33) Nureki, O; J Mol Biol 1994, V236, P710 HCAPLUS
- (34) Ohno, S; J Biochem 1998, V124, P1065 HCAPLUS
- (35) Ohno, S; J Biochem 2001, V130, P417 HCAPLUS
- (36) Scherberg, N; J Biol Chem 1978, V253, P1773 HCAPLUS
- (37) Sekine, S; J Biol Chem 2001, V276, P3723 HCAPLUS
- (38) Sharma, N; FEBS Lett 2000, V467, P37 HCAPLUS
- (39) Short, G; Biochemistry 1999, V38, P8808 HCAPLUS
- (40) Varshney, U; J Biol Chem 1991, V266, P24712 HCAPLUS
- (41) Wakasugi, K; EMBO J 1998, V17, P297 HCAPLUS
- (42) Wang, L; J Am Chem Soc 2000, V122, P5010 HCAPLUS
- (43) Wang, L; J Am Chem Soc 2002, V124, P1836 HCAPLUS
- (44) Wang, L; Science 2001, V292, P498 HCAPLUS
- (45) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS

L60 ANSWER 18 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:504945 HCAPLUS

DN 137:62163

TI Preparation of **recombinant protein** as chaperon  
**fusion protein**

IN Furutani, Masahiro; Hata, Junichi; Togi, Akiko

PA Sekisui Chemical Co., Ltd., Japan; Marine Biotechnology Institute Co.,  
Ltd.

SO PCT Int. Appl., 48 pp.

CODEN: PIXXD2

DT Patent

LA Japanese  
 IC ICM C12P021-02  
 ICS C12N015-09; C07K019-00  
 CC 15-2 (Immunochemistry)  
 Section cross-reference(s): 3

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002052029	A1	20020704	WO 2001-JP11438	20011226
	WO 2002052029	C2	20030424		
	W: AU, CA, JP, KR, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
PRAI	JP 2000-395740	A	20001226		
AB	<p>Provided is an <b>expression and cell-free translation</b> system for a <b>recombinant protein</b> as <b>fusion protein</b> with chaperonin subunit. A target <b>protein</b> is incorporated into the tertiary structure of chaperonin as a <b>fusion protein</b> with a chaperonin subunit (namely, a mol. chaperonin of about 60 kDa, heat shock <b>protein</b> of 60 kDa or thermosome) to thereby inhibit the <b>expression</b> of toxicity of the target <b>protein</b> in the host,. <b>Digestion</b> with a protease or CNBr at a methionine residue can be used to cleave off the target <b>protein</b> from the chaperonin subunit. Bacteria, archaea, or eukaryote derived chaperonin is used. Bacteria, yeast, animal <b>cell</b>, plant <b>cell</b>, insect <b>cell</b> can be used as host. Mammalian antibody heavy chain, light chain, single chain antibody Fv region, virus antigen, transmembrane receptor, or cytokine may be the target <b>protein</b>. <b>Expression</b> of hepatitis virus B surface antigen, hepatitis virus C core antigen, mouse anti-chicken lysozyme single chain antibody, human antibody heavy chain const. region, human interferon .alpha.2b, or human serotonin receptor as <b>fusion protein</b> with chaperonin GroEL, is described.</p>				
ST	<b>protein recombinant expression</b> chaperonin <b>fusion</b>				
IT	Chemokine receptors RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation) (CXCR4; prepn. of <b>recombinant protein</b> as chaperon <b>fusion protein</b> )				
IT	Chaperonins RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses) (GroEL; prepn. of <b>recombinant protein</b> as chaperon <b>fusion protein</b> )				
IT	Animal cell Bacteria (Eubacteria) <b>Escherichia coli</b> Insecta Plant cell Yeast (as host; prepn. of <b>recombinant protein</b> as chaperon <b>fusion protein</b> )				
IT	Archaeobacteria (Archaea) Eukaryota (chaperonin from; prepn. of <b>recombinant protein</b> as chaperon <b>fusion protein</b> )				
IT	Immunoglobulins RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation) (fragments, Fv region; prepn. of <b>recombinant protein</b> as chaperon <b>fusion protein</b> )				
IT	Immunoglobulins				

- RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(heavy chains; prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT Antigens  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(hepatitis B core; prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT Immunoglobulins  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(light chains; prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT 5-HT receptors  
Antibodies  
Antigens  
Cytokines  
**Fusion proteins (chimeric proteins)**  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT Chaperonins  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT Heat-shock **proteins**  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT Antibodies  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(single chain; prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT Antigens  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(surface; prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT Interferons  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(.alpha.-2b; prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT 439059-34-4, 1: PN: WO02052029 SEQID: 1 unclaimed DNA 439059-35-5, 2: PN: WO02052029 SEQID: 3 unclaimed DNA 439059-36-6, 3: PN: WO02052029 SEQID: 4 unclaimed DNA 439059-37-7, 4: PN: WO02052029 SEQID: 5 unclaimed DNA 439059-38-8, 5: PN: WO02052029 SEQID: 6 unclaimed DNA 439059-39-9, 6: PN: WO02052029 SEQID: 7 unclaimed DNA 439059-40-2, 7: PN: WO02052029 SEQID: 8 unclaimed DNA 439059-41-3, 8: PN: WO02052029 SEQID: 2 unclaimed DNA  
RL: PRP (Properties)  
(unclaimed nucleotide sequence; prepn. of **recombinant protein** as chaperon **fusion protein**)
- IT 9001-92-7, Proteinase  
RL: BUU (Biological use, unclassified); CAT (Catalyst use); BIOL (Biological study); USES (Uses)  
(use in cleavage; prepn. of **recombinant protein** as chaperon **fusion protein**)

IT 506-68-3, Cyanogen bromide ((CN)Br)  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(use in cleavage; prepn. of **recombinant protein** as  
chaperon **fusion protein**)

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE  
(1) Duenas, M; Biotechniques 1994, V16(3), P476 HCAPLUS  
(2) Korpela, T; WO 0066756 A1 2000 HCAPLUS  
(3) Korpela, T; EP 1173592 A1 2000 HCAPLUS  
(4) Nishihara, K; APPLIED AND ENVIRONMENTAL MICROBIOLOGY 1998, V64(5), P1694  
HCAPLUS

L60 ANSWER 19 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2002:442973 HCAPLUS  
DN 137:29466  
TI High-throughput **cell-free protein**  
**expression** system for structural genomics and proteomics studies  
AU Kigawa, Takanori; Yokoyama, Shigeyuki  
CS Genomic Sci. Cent., RIKEN, Japan  
SO Tanpakushitsu Kakusan Koso (2002), 47(8, Zokan), 1014-1019  
CODEN: TAKKAJ; ISSN: 0039-9450  
PB Kyoritsu Shuppan  
DT Journal; General Review  
LA Japanese  
CC 6-0 (General Biochemistry)  
Section cross-reference(s): 9  
AB A review on tech. development of high-throughput **protein**  
**expression** using **cell-free protein**  
**synthesis** system and its application to structure proteomics  
studies.  
ST review structure proteomics **protein expression** system  
IT **Translation, genetic**  
(**cell-free**; high-throughput **cell-**  
**free protein expression** system for  
structural genomics and proteomics studies)  
IT **Proteins**  
RL: BPN (Biosynthetic preparation); PRP (Properties); BIOL (Biological  
study); PREP (Preparation)  
(high-throughput **cell-free protein**  
**expression** system for structural genomics and proteomics  
studies)  
IT Genome  
(structural genomics; high-throughput **cell-free**  
**protein expression** system for structural genomics and  
proteomics studies)  
IT Proteome  
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL  
(Biological study)  
(structure proteomics; high-throughput **cell-free**  
**protein expression** system for structural genomics and  
proteomics studies)

L60 ANSWER 20 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2002:235434 HCAPLUS  
DN 137:164318  
TI **Expression of FLAG Fusion Proteins** in Insect  
Cells: Application to the Multi-subunit Transcription/DNA Repair Factor  
TFIIH  
AU Jawhari, Anass; Uhring, Muriel; Crucifix, Corinne; Fribourg, Sebastien;  
Schultz, Patrick; Poterszman, Arnaud; Egly, Jean Marc; Moras, Dino  
CS Institut de Genetique et de Biologie Cellulaire et Moleculaire,  
CNRS/INSERM/ULP, College de France, Illkirch, 67404, Fr.  
SO Protein Expression and Purification (2002), 24(3), 513-523

CODEN: PEXPEJ; ISSN: 1046-5928

PB Elsevier Science

DT Journal

LA English

CC 3-2 (Biochemical Genetics)

Section cross-reference(s): 12, 13

AB The multi-subunit transcription/DNA repair factor TFIIH was used as a model system to show that the **expression** of FLAG **fusion proteins** in insect cells constitutes a versatile tool for both structural and **functional** investigations. In the present study, we have constructed **recombinant** baculoviruses **expressing** the four core TFIIH subunits **fused** at their N-terminus to the FLAG peptide. Using these **recombinant** viruses we have established protocols based on anti-FLAG immunoaffinity chromatog. that allow the systematic anal. of pairwise interaction within multiprotein complexes and have developed a double tag strategy (FLAG and hexahistidine tags) for the identification and purifn. of stable TFIIH subcomplexes. A simple purifn. procedure was developed that leads to the isolation of **recombinant** TFIIH contg. the full set of subunits. The purified **recombinant** TFIIH was shown to be active in a transcription assay and to be structurally homologous to the endogenous complex by electron microscopy and image anal.

ST human multisubunit TFIIH **expression** purifn baculovirus vector Sf9 cell

IT **Enzymes**, preparation

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

(DNA helicase, XPB, His-tagged TFIIH subunit; **expression** of

FLAG **fusion proteins** in insect cells, application

to the human multi-subunit transcription/DNA repair factor TFIIH)

IT Animal cell line

(SF9, infection; **expression** of FLAG **fusion**

**proteins** in insect cells, application to the human

multi-subunit transcription/DNA repair factor TFIIH)

IT Transcription factors

RL: PUR (Purification or recovery); PREP (Preparation)

(TFIIH (transcription factor IIH); **expression** of FLAG

**fusion proteins** in insect cells, application to the

human multi-subunit transcription/DNA repair factor TFIIH)

IT Transcription, genetic

(cell-free; **expression** of FLAG

**fusion proteins** in insect cells,

application to the human multi-subunit transcription/DNA repair factor TFIIH)

IT Autographa californica nucleopolyhedrovirus

Human

Molecular **cloning**

(**expression** of FLAG **fusion proteins** in

insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)

IT **Fusion proteins (chimeric proteins)**

RL: BPN (Biosynthetic preparation); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation)

(**expression** of FLAG **fusion proteins** in

insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)

IT Molecular association

(of TFIIH subunits; **expression** of FLAG **fusion**

**proteins** in insect cells, application to the human

multi-subunit transcription/DNA repair factor TFIIH)

IT **Proteins**

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

- (p34, FLAG-tagged TFIIH subunit; **expression** of FLAG **fusion proteins** in insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)
- IT **Proteins**  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (p44, FLAG-tagged TFIIH subunit; **expression** of FLAG **fusion proteins** in insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)
- IT **Proteins**  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (p52, FLAG-tagged TFIIH subunit; **expression** of FLAG **fusion proteins** in insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)
- IT **Proteins**  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (p62, FLAG-tagged TFIIH subunit; **expression** of FLAG **fusion proteins** in insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)
- IT **Viral vectors**  
 (pSK277; **expression** of FLAG **fusion proteins** in insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)
- IT **Simulation and Modeling, physicochemical**  
 (**three-dimensional**; **expression** of FLAG **fusion proteins** in insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)
- IT 64134-30-1, Hexahistidine 98849-88-8, FLAG peptide  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (**expression** of FLAG **fusion proteins** in insect cells, application to the human multi-subunit transcription/DNA repair factor TFIIH)

RE.CNT 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Coin, F; EMBO J 1999, V18, P1357 HCAPLUS
- (2) Coin, F; Nature Genet 1998, V20, P184 HCAPLUS
- (3) Fukuda, A; Genes Cells 2001, V6, P707 HCAPLUS
- (4) Gerard, M; J Biol Chem 1991, V266, P20940 HCAPLUS
- (5) Hoeijmakers, J; Curr Opin Genet Dev 1996, V6, P26 HCAPLUS
- (6) Iyer, N; Biochemistry 1996, V35, P2157 HCAPLUS
- (7) Jarvik, J; Annu Rev Genet 1998, V32, P601 HCAPLUS
- (8) Kershner, E; J Biol Chem 1998, V273, P34444 HCAPLUS
- (9) Koh, S; Biotechniques 1997, V23, P622 HCAPLUS
- (10) Le Roy, G; J Biol Chem 1998, V273, P7134 HCAPLUS
- (11) Reardon, J; Proc Natl Acad Sci USA 1996, V93, P6482 HCAPLUS
- (12) Reardon, J; published erratum appears in Proc Natl Acad Sci USA 1996, V93(19), P10538 HCAPLUS
- (13) Schultz, P; Cell 2000, V102, P599 HCAPLUS
- (14) Svejstrup, J; Trends Biochem Sci 1996, V21, P346 HCAPLUS
- (15) Tirode, F; Mol Cell 1999, V3, P87 HCAPLUS
- (16) van Heel, M; J Struct Biol 1996, V116, P17 MEDLINE
- (17) Watanabe, Y; Genes Cells 2000, V5, P407 HCAPLUS
- (18) Winkler, G; J Biol Chem 1998, V273, P1092 HCAPLUS

L60 ANSWER 21 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2002:123933 HCAPLUS

DN 137:88980

TI An unnatural base pair for incorporating amino acid analogs into **proteins**

AU Hirao, Ichiro; Ohtsuki, Takashi; Fujiwara, Tsuyoshi; Mitsui, Tsuneo;

Yokogawa, Tomoko; Okuni, Taeko; Nakayama, Hiroshi; Takio, Koji; Yabuki, Takashi; **Kigawa, Takanori**; Kodama, Koichiro; Yokogawa, Takashi; Nishikawa, Kazuya; **Yokoyama, Shigeyuki**

CS Yokoyama Cytologic Project, ERATO, JST, c/o RIKEN, Wako-shi, Saitama, 351-0198, Japan

SO Nature Biotechnology (2002), 20(2), 177-182

CODEN: NABIF9; ISSN: 1087-0156

PB Nature America Inc.

DT Journal

LA English

CC 3-1 (Biochemical Genetics)

Section cross-reference(s): 6, 9

AB An unnatural base pair of 2-amino-6-(2-thienyl)purine (denoted by s) and pyridin-2-one (denoted by y) was developed to expand the genetic code. The ribonucleoside triphosphate of y was site-specifically incorporated into RNA, opposite s in a template, by T7 RNA polymerase. This transcription was coupled with translation in an **Escherichia coli cell-free** system. The yAG codon in the transcribed ras mRNA was recognized by the CUs anticodon of a yeast tyrosine tRNA (tRNA) variant, which had been **enzymically** aminoacylated with an unnatural amino acid, 3-chlorotyrosine. Site-specific incorporation of 3-chlorotyrosine into the Ras **protein** was demonstrated by liq. chromatog.-mass spectrometry (LC-MS) anal. of the products. This coupled transcription-translation system will permit the efficient **synthesis of proteins** with a tyrosine analog at the desired position.

ST **protein** engineering chlorotyrosine genetic translation unnatural base pair; tyrosine tRNA RNA sequence human gene cHa ras **protein**

IT Amino acids, preparation

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

(analog; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)

IT Nucleic acid bases

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(analog; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)

IT Gene, animal

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(c-Ha-ras, mRNA, yAG codon-contg.; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)

IT Transcription, genetic

**Translation, genetic**

(**cell-free**; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)

IT DNA

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(double-stranded, template, s .cntdot. T base pair; unnatural base pair of 2-amino-6-(2-thienyl)purine (denoted s) and pyridin-2-one (denoted y) for incorporating amino acid analogs into **proteins** by a coupled **cell-free** transcription-translation reaction)

IT **mRNA sequences**

(of human c-Ha-ras with incorporated pyridin-2-one (y))

IT Ras **proteins**

RL: BPN (Biosynthetic preparation); PRP (Properties); BIOL (Biological



- study); PREP (Preparation)  
(p21c-Ha-ras, 3-chlorotyrosine-modified; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT Nucleic acid bases  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(pairing; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT Anticodons  
RL: BUU (Biological use, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(**synthetic**, CUs; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT Codons  
RL: BUU (Biological use, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(**synthetic**, yAG; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT tRNA  
RL: BPN (Biosynthetic preparation); BUU (Biological use, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(tyrosine-specific, CUs anticodon-contg.; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT Human  
**Protein** engineering  
(unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT RNA  
RL: BPN (Biosynthetic preparation); PRP (Properties); BIOL (Biological study); PREP (Preparation)  
(unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT 440200-36-2P  
RL: BPN (Biosynthetic preparation); PRP (Properties); BIOL (Biological study); PREP (Preparation)  
(amino acid sequence; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT 439618-75-4P  
RL: BPN (Biosynthetic preparation); BUU (Biological use, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(nucleotide sequence; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT 439618-71-0P  
RL: BPN (Biosynthetic preparation); PRP (Properties); BIOL (Biological study); PREP (Preparation)  
(nucleotide sequence; unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with 3-chlorotyrosine at position 32)
- IT 7423-93-0P  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(unnatural base pair for incorporating amino acid analogs into **proteins**, prodn. of human Ras **protein** with

3-chlorotyrosine at position 32)  
IT 142-08-5, Pyridin-2-one 156489-35-9 382146-88-5  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)

(unnatural base pair of 2-amino-6-(2-thienyl)purine (denoted s) and  
pyridin-2-one (denoted y) for incorporating amino acid analogs into  
**proteins** by a coupled **cell-free**  
transcription-translation reaction)

RE.CNT 52 THERE ARE 52 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Bain, J; J Am Chem Soc 1989, V111, P8013 HCAPLUS
- (2) Bain, J; Nature 1992, V356, P537 HCAPLUS
- (3) Benner, S; The RNA world Edn 2 1999, P163 HCAPLUS
- (4) Boeck, A; Science 2001, V292, P453
- (5) Chow, C; J Biol Chem 1993, V268, P12855 HCAPLUS
- (6) Cload, S; Chem Biol 1998, V3, P1033
- (7) Cornish, V; Proc Natl Acad Sci 1994, V91, P2910 HCAPLUS
- (8) Doering, V; Science 2001, V292, P5001
- (9) Fujiwara, T; Bioorg Med Chem Lett 2001, V11, P2221 HCAPLUS
- (10) Hecht, S; Acc Chem Res 1992, V25, P545 HCAPLUS
- (11) Hocek, M; Collect Czech Chem Commun 2001, V66, P483 HCAPLUS
- (12) Hosaka, T; J Am Chem Soc 1999, V121, P34
- (13) Ishikawa, M; Tetrahedron Lett 2000, V41, P3931 HCAPLUS
- (14) Judice, J; Science 1993, V261, P1578 HCAPLUS
- (15) Kawasaki, H; Anal Biochem 1990, V191, P332 HCAPLUS
- (16) Kern, D; Eur J Biochem 1972, V31, P148 HCAPLUS
- (17) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (18) Kigawa, T; J Biomol NMR 1995, V6, P129 HCAPLUS
- (19) Kowal, A; Proc Natl Acad Sci 2001, V98, P2268 HCAPLUS
- (20) Leinfelder, W; Nature 1988, V331, P723 HCAPLUS
- (21) Lutz, M; Bioorg Med Chem Lett 1998, V8, P499 HCAPLUS
- (22) Madin, K; Proc Natl Acad Sci 2000, V97, P559 HCAPLUS
- (23) Magliery, T; J Mol Biol 2001, V307, P755 HCAPLUS
- (24) Mamaev, S; J Am Chem Soc 1996, V118, P7243 HCAPLUS
- (25) Masaki, T; Biochim Biophys Acta 1981, V650, P44
- (26) Matray, T; Nature 1999, V399, P704 HCAPLUS
- (27) Moore, B; J Mol Biol 2000, V298, P195 HCAPLUS
- (28) Morales, J; Nat Struct Biol 1998, V5, P950 HCAPLUS
- (29) Murakami, H; J Am Chem Soc 1998, V120, P7520 HCAPLUS
- (30) Noren, C; Science 1989, V244, P182 HCAPLUS
- (31) Nowak, M; Methods Enzymol 1998, V293, P504 HCAPLUS
- (32) Nowak, M; Science 1995, V268, P439 HCAPLUS
- (33) Ogawa, A; J Am Chem Soc 2000, V122, P3274 HCAPLUS
- (34) Ohno, S; J Biochem 1998, V124, P1065 HCAPLUS
- (35) Ohno, S; J Biochem 2001, V130, P417 HCAPLUS
- (36) Ohtsuki, T; Nucleic Acids Res 1996, V24, P662 HCAPLUS
- (37) Ohtsuki, T; Proc Natl Acad Sci 2001, V98, P4922 HCAPLUS
- (38) Ohyama, T; J Biochem 1985, V97, P29 HCAPLUS
- (39) Piccirilli, J; Nature 1990, V343, P33 HCAPLUS
- (40) Rappaport, H; Biochemistry 1993, V32, P3047 HCAPLUS
- (41) Saks, M; J Biol Chem 1996, V271, P23169 HCAPLUS
- (42) Service, R; Science 2000, V289, P232 HCAPLUS
- (43) Shimizu, Y; Nat Biotechnol 2001, V19, P751 HCAPLUS
- (44) Short, G; Biochemistry 1999, V38, P8808 HCAPLUS
- (45) Short, G; Biochemistry 2000, V39, P8768 HCAPLUS
- (46) Spirin, A; Science 1988, V242, P1162 HCAPLUS
- (47) Switzer, C; Biochemistry 1993, V32, P10489 HCAPLUS
- (48) Turcatti, G; J Biol Chem 1996, V271, P19991 HCAPLUS
- (49) van Hest, J; J Am Chem Soc 2000, V122, P1282 HCAPLUS
- (50) Wang, L; Science 2001, V292, P498 HCAPLUS
- (51) Wu, Y; J Am Chem Soc 2000, V122, P7621 HCAPLUS
- (52) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS

L60 ANSWER 22 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2002:47341 HCAPLUS  
DN 137:104208  
TI The many faces of **protein expression**  
AU Janssen, Deborah  
CS USA  
SO Genomics & Proteomics (2001), 1(3), 53-57  
CODEN: GPERA6; ISSN: 1536-7495  
PB Cahners Business Information  
DT Journal; General Review  
LA English  
CC 3-0 (Biochemical Genetics)  
Section cross-reference(s): 6, 16  
AB A review. Various **protein expression** systems that provide easy, rapid, accurate **protein** characterization for a wide range of user application have been developed. One such system is called pET (plasmid for **expression** by T7 RNA polymerase) **expression** system, developed by Novagen Inc., for eukaryotic genes in **E. coli**. The Novagen pET **expression** system offers 39 different pET **vectors**, 36 different **expression** strains, 14 **fusion** tag choices, and many options for rapid anal. and purifn. of target **proteins**. Another system is the Rapid Translation System (RTS), a **cell-free protein expression** platform, which eliminates the need for housekeeping cellular activities.  
ST review **protein expression** system bacterium eukaryote Escherichia  
IT **Escherichia coli**  
(**expression** host; com. **expression** systems for manuf. of **proteins** in prokaryotic and eukaryotic systems)  
IT **Translation, genetic**  
(in vitro, for **protein** manuf.; com. **expression** systems for manuf. of **proteins** in prokaryotic and eukaryotic systems)  
IT **Plasmid vectors**  
(pET; com. **expression** systems for manuf. of **proteins** in prokaryotic and eukaryotic systems)  
IT **Proteins**  
RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
(prepn. by **translation** of; com. **expression** systems for manuf. of **proteins** in prokaryotic and eukaryotic systems)  
IT **Fermentation**  
(**protein**; com. **expression** systems for manuf. of **proteins** in prokaryotic and eukaryotic systems)

L60 ANSWER 23 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2001:586343 HCAPLUS  
DN 136:196343  
TI The **cell-free protein** biosynthesis - applications and analysis of the system  
AU Lamla, Thorsten; Mammeri, Kerstin; Erdmann, Volker A.  
CS Institut fur Biochemie, Freie Universitat Berlin, Berlin, D-14195, Germany  
SO Acta Biochimica Polonica (2001), 48(2), 453-465  
CODEN: ABPLAF; ISSN: 0001-527X  
PB Polish Biochemical Society  
DT Journal  
LA English  
CC 9-2 (Biochemical Methods)  
Section cross-reference(s): 6  
AB The in vitro **protein** biosynthesis has the potentials to become a powerful technol. for biochem. research. Beside the detn. of structure and **function** the in vitro evolution of **proteins** is

also of great interest. The system described was used to produce bovine heart fatty acid binding **protein** (FABP) and bacterial chloramphenicol acetyltransferase (CAT) with and without **fusion** of the Strep-tag II affinity peptide. The **proteins** were purified after and during **protein** biosynthesis by using a StrepTactin Sepharose matrix. No significant influence of the Strep-tag and the conditions during the affinity chromatog. on maturation or activity of the **protein** was obsd. The in vitro evolution of **proteins** is feasible by means of ribosome display. The selection of a specific mRNA coding for a shortened FABP with a N-terminal His-tag via the accompanying **protein** property was shown. Goal of the selection was to bind the FABP via the His-tag on Ni(II)-IDA-agarose. After nine cycles of transcription, translation, affinity selection and RT-PCR the **protein** with the His-tag could be enriched 108-fold. In order to correlate a possible relationship between changes in **protein** population and biol. **function** studies were initiated in which 2-dimensional **protein** patterns of the total in vitro system were compared after 0 and 2 h reaction time. The very interesting findings are that a no. of **proteins** disappear, while others are newly formed during **protein synthesis**.

- ST **cell free** system **protein** biosynthesis; fatty acid binding **protein** biosynthesis; chloramphenicol acetyltransferase biosynthesis
- IT **Proteins**  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (FABP (fatty acid-binding **protein**); **cell-free protein** biosynthesis - applications and anal. of system)
- IT **Translation, genetic**  
 (cell-free **protein** biosynthesis - applications and anal. of system)
- IT mRNA  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (cell-free **protein** biosynthesis - applications and anal. of system)
- IT Gel electrophoresis  
 (two-dimensional; **cell-free protein** biosynthesis - applications and anal. of system)
- IT 9040-07-7P, Chloramphenicol acetyltransferase  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (cell-free **protein** biosynthesis - applications and anal. of system)

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE

- (1) Appel, R; Electrophoresis 1991, V12, P722 HCAPLUS
- (2) Bradford, M; Anal Biochem 1976, V72, P248 HCAPLUS
- (3) Cronenberger, J; J Mol Biol 1975, V95, P125 HCAPLUS
- (4) Hanes, J; Proc Natl Acad Sci USA 1997, V94, P4937 HCAPLUS
- (5) Haukanes, B; Biotechnology 1993, V11, P60 MEDLINE
- (6) Heukeshoven, J; Electrophoresis 1985, V6, P103 HCAPLUS
- (7) Holschuh, K; J Biol Chem 1982, V257, P1987 HCAPLUS
- (8) Klose, J; Electrophoresis 1995, V16, P1034 HCAPLUS
- (9) Laemmli, U; Nature 1970, V227, P680 HCAPLUS
- (10) Mattheakis, L; Proc Natl Acad Sci USA 1994, V91, P9022 HCAPLUS
- (11) Merk, H; J Biochem 1999, V125, P328 HCAPLUS
- (12) Moore, J; Nat Biotechnol 1996, V14, P458 HCAPLUS
- (13) Nygren, P; Trends Biotechnol 1994, V12, P184 HCAPLUS
- (14) Osborne, S; Chem Rev 1997, V97, P349 HCAPLUS
- (15) Peled-Zehavi, H; Methods Enzymol 2000, V318, P297 HCAPLUS
- (16) Roberts, R; Proc Natl Acad Sci USA 1997, V94, P12297 HCAPLUS

- (17) Ryabova, L; Nucleic Acids Res 1989, V17, P4412 HCAPLUS
- (18) Sambrook, J; Molecular cloning 1989
- (19) Schatz, P; Methods Enzymol 1996, V267, P171 HCAPLUS
- (20) Schindler, P; Electrophoresis 1999, V20, P806 HCAPLUS
- (21) Schmidt, T; J Mol Biol 1996, V255, P753 HCAPLUS
- (22) Siemann, M; Eur J Biochem, in press 2000
- (23) Smith, G; Chem Rev 1997, V97, P391 HCAPLUS
- (24) Spirin, A; Science 1988, V242, P1162 HCAPLUS
- (25) Stiege, W; J Biotechnol 1995, V41, P81 HCAPLUS
- (26) Triana-Alonso, F; J Biol Chem 1995, V270, P6298 HCAPLUS
- (27) Voss, S; Protein Eng 1997, V10, P975 HCAPLUS
- (28) Zhang, J; Proc Natl Acad Sci USA 1997, V94, P4504 HCAPLUS

L60 ANSWER 24 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2001:232988 HCAPLUS

DN 135:41519

TI **Cell-Free** Production of Biologically Active  
Polypeptides: Application to the **Synthesis** of Antibacterial  
Peptide Cecropin

AU Martemyanov, Kirill A.; Shirokov, Vladimir A.; Kurnasov, Oleg V.; Gudkov,  
Anatoly T.; Spirin, Alexander S.

CS Institute of Protein Research, Russian Academy of Sciences, Pushchino,  
Moscow Region, 142292, Russia

SO Protein Expression and Purification (2001), 21(3), 456-461  
CODEN: PEXPEJ; ISSN: 1046-5928

PB Academic Press

DT Journal

LA English

CC 3-1 (Biochemical Genetics)  
Section cross-reference(s): 6

AB An approach to preparative prodn. of polypeptides, including uneasily  
testable, degradable, and toxic ones, is proposed on the basis of in vitro  
**expression** systems of last generation, such as continuous-exchange  
**cell-free** and continuous-flow **cell-**  
**free** transcription-translation systems. The approach implies that  
a polypeptide of interest is **synthesized** as a **fusion**  
**protein** with the polypeptide linked to **green**  
**fluorescent protein (GFP)** through a cleavable  
spacer. The **GFP** moiety provides direct visualization and quant.  
monitoring of the polypeptide **synthesis**, as well as **soly**  
. and stability of the product. The **synthesis** of  
**functionally** active antibacterial polypeptide cecropin P1 (31  
amino acid residues) has been demonstrated. (c) 2001 Academic Press.

ST toxic **protein fusion** product **cell**  
**free synthesis**; labile **protein fusion**  
product **cell free synthesis**; **green**  
**fluorescent protein fusion** product  
**cell free synthesis**; translation in vitro  
toxic labile **protein fusion** product; cecropin manuf  
**fusion** product **cell free** translation

IT Gene, microbial

RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(10, ribosome binding site of; **synthesis** of toxic or labile  
**proteins** as cleavable **fusion** products with  
**green fluorescent protein** in **cell**  
**-free** systems)

IT Transcription, genetic  
(**cell-free**, translation coupled; **synthesis**  
of toxic or labile **proteins** as cleavable **fusion**  
products with **green fluorescent protein**  
in **cell-free** systems)

IT Translation, genetic  
(**cell-free**; **synthesis** of toxic or labile

**proteins as cleavable fusion products with  
green fluorescent protein in cell  
-free systems)**

IT. **Proteins, specific or class**

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

**(green fluorescent, fusion  
products; synthesis of toxic or labile  
proteins as cleavable fusion products with  
green fluorescent protein in cell  
-free systems)**

IT Genetic element

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

**(ribosome-binding site, gene 10, in transcription translation  
constructs; synthesis of toxic or labile proteins  
as cleavable fusion products with green  
fluorescent protein in cell-free  
systems)**

IT Coliphage T7

**(transcription-translation vectors derived from;  
synthesis of toxic or labile proteins as cleavable  
fusion products with green fluorescent  
protein in cell-free systems)**

IT 506-68-3, Cyanogen bromide

RL: RCT (Reactant); RACT (Reactant or reagent)  
**(fusion proteins cleavable by; synthesis  
of toxic or labile proteins as cleavable fusion  
products with green fluorescent protein  
in cell-free systems)**

IT 9014-74-8, Enterokinase

RL: CAT (Catalyst use); USES (Uses)  
**(fusion proteins contg. cleavage site for;  
synthesis of toxic or labile proteins as cleavable  
fusion products with green fluorescent  
protein in cell-free systems)**

IT 80802-79-5DP, Cecropin, **fusion products with green  
fluorescent protein**

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

**(synthesis of toxic or labile proteins as cleavable  
fusion products with green fluorescent  
protein in cell-free systems)**

RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Alakhov, Y; US 5478730 1988 HCAPLUS
- (2) Baranov, V; Gene 1989, V84, P463 HCAPLUS
- (3) Baranov, V; Methods Enzymol 1993, V217, P123 HCAPLUS
- (4) Buchner, J; Biotechnology 1992, V10, P682 HCAPLUS
- (5) Chalfie, M; Science 1994, V263, P802 HCAPLUS
- (6) Chekulayeva, M; Continuous-exchange cell-free protein-synthesizing system:  
Synthesis of HIV antigen Nef fused with green fluorescent protein  
Submitted for publication 1999
- (7) Cormack, B; Gene 1996, V173, P33 HCAPLUS
- (8) Cramer, A; Nature Biotechnol 1996, V14, P315 HCAPLUS
- (9) Hancock, R; Trends Biotechnol 1998, V16, P82 HCAPLUS
- (10) Hara, S; Biochem Biophys Res Commun 1996, V220, P664 HCAPLUS
- (11) Haught, C; Biotech Bioeng 1998, V57, P55 HCAPLUS
- (12) Jermutus, L; Curr Opin Biotechnol 1998, V9, P534 HCAPLUS
- (13) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (14) Kim, D; Biotechnol Prog 1996, V12, P645 HCAPLUS
- (15) Laemmli, U; Nature 1970, V227, P680 HCAPLUS
- (16) Lehrer, R; J Immunol Methods 1991, V137, P167 HCAPLUS

- (17) Lepage, P; Anal Biochem 1993, V213, P40 HCAPLUS  
 (18) Martemyanov, K; Biotechnol Lett 1996, V18, P1357 HCAPLUS  
 (19) Martemyanov, K; FEBS Lett 1997, V414, P268 HCAPLUS  
 (20) Pang, S; Gene 1992, V116, P165 HCAPLUS  
 (21) Piers, K; Gene 1993, V134, P7 HCAPLUS  
 (22) Spirin, A; Frontiers of Bioprocessing 2 1992, P31 HCAPLUS  
 (23) Spirin, A; Science 1988, V242, P1162 HCAPLUS  
 (24) Wang, L; Biochem Mol Biol Int 1997, V41, P1051 HCAPLUS  
 (25) Xie, W; Biochem Mol Biol Int 1996, V39, P487 HCAPLUS  
 (26) Zubay, G; Annu Rev Genet 1973, V7, P267 HCAPLUS

L60 ANSWER 25 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2001:152863 HCAPLUS

DN 134:204756

TI Methods for the detection, analysis and isolation of nascent  
**proteins**

IN Rothschild, Kenneth J.; Gite, Sadanand; Olejnik, Jerzy

PA Ambergen, Inc., USA

SO PCT Int. Appl., 204 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C12P021-06

ICS C12N009-00

CC 9-16 (Biochemical Methods)

Section cross-reference(s): 1, 3

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001014578	A1	20010301	WO 2000-US23233	20000823
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	US 6303337	B1	20011016	US 1999-382950	19990825
	US 6306628	B1	20011023	US 1999-382736	19990825
	EP 1210449	A1	20020605	EP 2000-957758	20000823
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
	US 2002132248	A1	20020919	US 2001-973145	20011009
	US 2003092031	A1	20030515	US 2002-174368	20020618
PRAI	US 1999-382736	A	19990825		
	US 1999-382950	A	19990825		
	WO 2000-US23233	W	20000823		
	US 2002-49332	A2	20020621		
AB	This invention relates to non-radioactive markers that facilitate the detection and anal. of nascent <b>proteins translated</b> within cellular or <b>cell-free translation</b> systems. Nascent <b>proteins</b> contg. these markers can be rapidly and efficiently detected, isolated and analyzed without the handling and disposal problems assocd. with radioactive reagents. Preferred markers are dipyrrometheneboron difluoride (4,4-difluoro-4-bora-3a,4a-diaza-s-indacene) dyes.				
ST	detection analysis isolation nascent <b>protein</b>				
IT	Codons				
	RL: BSU (Biological study, unclassified); BIOL (Biological study) (AUG; methods for detection, anal. and isolation of nascent <b>proteins</b> )				

- IT **Proteins**, specific or class  
RL: ANT (Analyte); ANST (Analytical study)  
(Immunogenic; methods for detection, anal. and isolation of nascent **proteins**)
- IT **Proteins**, specific or class  
RL: ANT (Analyte); ANST (Analytical study)  
(**Recombinant**; methods for detection, anal. and isolation of nascent **proteins**)
- IT Wheat  
(germ; methods for detection, anal. and isolation of nascent **proteins**)
- IT Transformation, neoplastic  
(immortalization; methods for detection, anal. and isolation of nascent **proteins**)
- IT tRNA  
RL: BUU (Biological use, unclassified); RCT (Reactant); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)  
(initiator; methods for detection, anal. and isolation of nascent **proteins**)
- IT **Proteins**, specific or class  
RL: ANT (Analyte); ANST (Analytical study)  
(lipid-binding; methods for detection, anal. and isolation of nascent **proteins**)
- IT Affinity  
Aminoacylation  
Animal tissue culture  
Bacteria (Eubacteria)  
Cell  
Dialysis  
Dog (Canis familiaris)  
Drugs  
Dyes  
**Escherichia coli**  
Flow  
Fluorometry  
Frog  
Immunoassay  
Insect (Insecta)  
Mutation  
Neoplasm  
PCR (polymerase chain reaction)  
Pancreas  
Parasite  
Rabbit  
Reticulocyte  
Temperature effects, biological  
Test kits  
**Translation, genetic**  
Virus  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT Carbohydrates, analysis  
Cytokines  
**Enzymes**, analysis  
**Fusion proteins (chimeric proteins)**  
Hormones, animal, analysis  
**Protein** formation factors  
RL: ANT (Analyte); ANST (Analytical study)  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT Proteome  
RL: ANT (Analyte); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study)



- (methods for detection, anal. and isolation of nascent **proteins**)
- IT Gene, animal  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT Primers (nucleic acid)  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT tRNA  
RL: BUU (Biological use, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(misaminoacylated; methods for detection, anal. and isolation of nascent **proteins**)
- IT **Proteins**, general, analysis  
RL: ANT (Analyte); PUR (Purification or recovery); ANST (Analytical study); PREP (Preparation)  
(nascent; methods for detection, anal. and isolation of nascent **proteins**)
- IT **Proteins**, specific or class  
RL: ANT (Analyte); ANST (Analytical study)  
(nucleic acid-binding; methods for detection, anal. and isolation of nascent **proteins**)
- IT Egg  
(oocyte; methods for detection, anal. and isolation of nascent **proteins**)
- IT tRNA  
RL: BUU (Biological use, unclassified); RCT (Reactant); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)  
(suppressor; methods for detection, anal. and isolation of nascent **proteins**)
- IT Hemolysins  
RL: ANT (Analyte); BPR (Biological process); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); PROC (Process)  
(.alpha.-; methods for detection, anal. and isolation of nascent **proteins**)
- IT 9001-92-7, Protease  
RL: ANT (Analyte); ANST (Analytical study)  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT 138026-71-8, Bodipy  
RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); RCT (Reactant); ANST (Analytical study); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT 91-64-5DP, Coumarin, derivs.  
RL: ARG (Analytical reagent use); RCT (Reactant); SPN (Synthetic preparation); ANST (Analytical study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT 71-00-1, Histidine, biological studies  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(methods for detection, anal. and isolation of nascent **proteins**)
- IT 60-32-2, 6-Aminocaproic acid 3992-42-5 35231-44-8 58178-21-5 69976-70-1 87424-19-9 105047-45-8  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(methods for detection, anal. and isolation of nascent **proteins**)

)

IT 58-61-7P, Adenosine, preparation 610-15-1P 6851-99-6P 35013-72-0P  
 72040-64-3P 87424-17-7P 110744-77-9P 133852-19-4P 133852-20-7P  
 174406-74-7P 208660-68-8P 250610-57-2P 328387-23-1P 328387-40-2P  
 328387-48-0P 328387-50-4P 328387-52-6P 328387-54-8P 328387-56-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (methods for detection, anal. and isolation of nascent **proteins**)

)

IT 133852-21-8P 328387-26-4P 328387-45-7DP, reaction with amino acids  
 328564-07-4P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (methods for detection, anal. and isolation of nascent **proteins**)

)

IT 328601-46-3 328601-47-4, 2: PN: WO0114578 PAGE: 15 unclaimed RNA  
 328601-48-5, 3: PN: WO0114578 PAGE: 15 unclaimed RNA  
 RL: PRP (Properties)  
 (unclaimed nucleotide sequence; methods for the detection, anal. and  
 isolation of nascent **proteins**)

IT 64134-30-1 92000-76-5 98849-88-8 103425-05-4 112218-36-7  
 145646-22-6 169802-74-8 169802-75-9 174144-06-0 205938-74-5  
 245445-88-9 268741-28-2 328547-57-5 328547-58-6 328601-49-6  
 328601-50-9 328601-51-0 328601-52-1 328601-53-2 328601-54-3  
 328601-55-4 328601-56-5 328601-57-6 328601-58-7 328601-59-8  
 328601-60-1 328601-61-2 328601-62-3 328601-63-4 328601-64-5  
 328601-65-6 328601-66-7  
 RL: PRP (Properties)  
 (unclaimed sequence; methods for the detection, anal. and isolation of  
 nascent **proteins**)

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Kang; US 5451663 A 1995 HCAPLUS  
 (2) Kudlicki; J Mol Biol 1994, V244, P319 HCAPLUS  
 (3) Rothschild; US 5643722 A 1997 HCAPLUS  
 (4) Turcatti; J Biol Chem 1996, V271(33), P19991 HCAPLUS

L60 ANSWER 26 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2001:84778 HCAPLUS

DN 135:149273

TI Structure determination of **protein folds** using the  
**cell-free synthesis** and NMR spectroscopy

AU Kigawa, Takanori; Yokoyama, Shigeyuki

CS Genomic Sciences Center, RIKEN, Japan

SO Jikken Igaku (2000), 18(18), 2464-2468  
 CODEN: JIIGEF; ISSN: 0288-5514

PB Yodosha

DT Journal; General Review

LA Japanese

CC 9-0 (Biochemical Methods)  
 Section cross-reference(s): 6

AB A review with 11 refs. on automation and fast detn. of **protein**  
 structure, **protein folding**, and physiol.  
**function(s)** using **cell-free protein**  
**synthesis** system to manuf. target **protein**, and NMR to  
 study the structure of the target **protein**.

ST review **protein** structure detn automation

IT **Secondary structure**  
**Tertiary structure**  
 (protein; structure detn. of **protein folds**  
 using **cell-free synthesis** and NMR  
 spectroscopy)

IT **Protein folding**  
 (structure detn. of **protein folds** using

## cell-free synthesis and NMR spectroscopy)

L60 ANSWER 27 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
 AN 2001:76271 HCAPLUS  
 DN 134:219310  
 TI Automated search of natively **folded protein** fragments  
 for high-throughput structure determination in structural genomics  
 AU Kuroda, Yutaka; Tani, Kazutoshi; Matsuo, Yo; **Yokoyama, Shigeyuki**  
 CS Protein Research Group, Genomic Sciences Center (GSC), The Institute of  
 Physical and Chemical Research (RIKEN), Kanagawa, 230-0045, Japan  
 SO Protein Science (2000), 9(12), 2313-2321  
 CODEN: PRCIEI; ISSN: 0961-8368  
 PB Cambridge University Press  
 DT Journal  
 LA English  
 CC 9-16 (Biochemical Methods)  
 Section cross-reference(s): 6  
 AB Structural genomic projects envision almost routine **protein**  
 structure detns., which are currently imaginable only for small  
**proteins** with mol. wts. below 25,000 Da. For larger  
**proteins**, structural insight can be obtained by breaking them into  
 small segments of amino acid sequences that can **fold** into native  
 structures, even when isolated from the rest of the **protein**.  
 Such segments are autonomously **folding** units (AFU) and have  
 sizes suitable for fast structural analyses. Here, we propose to expand  
 an intuitive procedure often employed for identifying biol. important  
**domains** to an automatic method for detecting putative  
**folded protein** fragments. The procedure is based on the  
 recognition that large **proteins** can be regarded as a combination  
 of independent **domains** conserved among diverse organisms. We  
 thus have developed a program that reorganizes the output of BLAST  
 searches and detects regions with a large no. of similar sequences. To  
 automate the detection process, it is reduced to a simple geometrical  
 problem of recognizing rectangular shaped elevations in a graph that plots  
 the no. of similar sequences at each residue of a query sequence. We used  
 our program to quant. corroborate the premise that segments with conserved  
 sequences correspond to **domains** that **fold** into native  
 structures. We applied our program to a test data set composed of 99  
 amino acid sequences contg. 150 segments with structures listed in the  
**Protein** Data Bank, and thus known to **fold** into native  
 structures. Overall, the fragments identified by our program have an  
 almost 50% probability of forming a native structure, and comparable  
 results are obsd. with sequences contg. **domain** linkers  
 classified in SCOP. Furthermore, we verified that our program identifies  
 AFU in libraries from various organisms, and we found a significant no. of  
 AFU candidates for structural anal., covering an estd. 5 to 20% of the  
 genomic databases. Altogether, these results argue that methods based on  
 sequence similarity can be useful for dissecting large **proteins**  
 into small autonomously **folding domains**, and such  
 methods may provide an efficient support to structural genomics projects.  
 ST **protein** conformation **folding** sequence homol program  
 IT Computer program  
 (PASS; automated search of natively **folded protein**  
 fragments for high-throughput structure detn. in structural genomics)  
 IT **Protein folding**  
**Protein** sequences  
 (automated search of natively **folded protein**  
 fragments for high-throughput structure detn. in structural genomics)  
 IT Peptides, analysis  
**Proteins**, general, analysis  
 RL: ANT (Analyte); BSU (Biological study, unclassified); PRP (Properties);  
 ANST (Analytical study); BIOL (Biological study)  
 (automated search of natively **folded protein**

fragments for high-throughput structure detn. in structural genomics)  
IT **Protein sequences**  
(homol.; automated search of natively **folded protein**  
fragments for high-throughput structure detn. in structural genomics)  
IT **Conformation**  
(**protein**; automated search of natively **folded**  
**protein** fragments for high-throughput structure detn. in  
structural genomics)

RE.CNT 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Altschul, S; J Mol Biol 1990, V215, P403 HCAPLUS
- (2) Altschul, S; Nucleic Acids Res 1997, V25, P3389 HCAPLUS
- (3) Bairoch, A; Nucleic Acids Res 1996, V24, P21 HCAPLUS
- (4) Baron, M; Nature 1990, V345, P642 HCAPLUS
- (5) Benson, D; Nucleic Acids Res 2000, V28, P15 HCAPLUS
- (6) Brenner, S; Nucleic Acids Res 1999, V27, P251 HCAPLUS
- (7) Burley, S; Nat Genet 1999, V23, P151 HCAPLUS
- (8) Campbell, I; Trends Biotechnol 1994, V12, P168 HCAPLUS
- (9) Clore, G; Nat Struct Biol Suppl 1997, V4, P849 HCAPLUS
- (10) Corpet, F; Nucleic Acids Res 1998, V26, P323 HCAPLUS
- (11) Elofsson, A; Bioinformatics 1999, V15, P480 HCAPLUS
- (12) Gaasterland, T; Nat Biotechnol 1998, V16, P625 HCAPLUS
- (13) Gomi, S; J Gen Virol 1999, V80, P1323 HCAPLUS
- (14) Gracy, J; Bioinformatics 1998, V14, P164 HCAPLUS
- (15) Henikoff, S; Bioinformatics 1999, V15, P471 HCAPLUS
- (16) Holm, L; Nucleic Acids Res 1998, V26, P316 HCAPLUS
- (17) Hoseki, J; J Biochem (Tokyo) 1999, V126, P951 HCAPLUS
- (18) Hubbard, S; Biochim Biophys Acta 1998, V1382, P191 HCAPLUS
- (19) Hubbard, T; Nucleic Acids Res 1999, V27, P254 HCAPLUS
- (20) Kabsch, W; Biopolymers 1983, V22, P2577 HCAPLUS
- (21) Kim, S; Nat Struct Biol Suppl 1998, V5, P643 HCAPLUS
- (22) Kohda, D; Cell 1993, V72, P953 HCAPLUS
- (23) Kohda, D; Cell 1996, V86, P767 HCAPLUS
- (24) Koshiha, S; FEBS Lett 1999, V442, P138 HCAPLUS
- (25) Kraulis, P; J Appl Crystallogr 1991, V1991, P946
- (26) Lugovskoy, A; Cell 1999, V99, P747 HCAPLUS
- (27) Mallick, P; Proc Natl Acad Sci USA 2000, V97, P2450 HCAPLUS
- (28) Murzin, A; J Mol Biol 1995, V247, P536 HCAPLUS
- (29) Nakagawa, N; J Biochem (Tokyo) 1999, V126, P986 HCAPLUS
- (30) Nicodeme, P; Bioinformatics 1998, V14, P508 HCAPLUS
- (31) Orengo, C; Curr Opin Struct Biol 1999, V9, P374 HCAPLUS
- (32) Orengo, C; Nature 1994, V372, P631 HCAPLUS
- (33) Orengo, C; Structure 1997, V5, P1093 HCAPLUS
- (34) Salzmann, M; Proc Natl Acad Sci USA 1998, V95, P13585 HCAPLUS
- (35) Shapiro, L; Structure 1998, V6, P265 HCAPLUS
- (36) Sonnhammer, E; Protein Sci 1994, V3, P482 HCAPLUS
- (37) Sonnhammer, E; Proteins 1997, V28, P405 HCAPLUS
- (38) Wuthrich, K; Nat Struct Biol Suppl 1998, V5, P492 HCAPLUS
- (39) Yokoyama, S; Nat Struct Biol 2000, V7(Suppl), P943
- (40) Yu, H; Science 1992, V258, P1665 HCAPLUS

L60 ANSWER 28 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2000:900304 HCAPLUS

DN 134:27258

TI Method and multi-channel membrane device for high throughput **cell**  
**free protein synthesis**

IN Metzler, Thomas; Schels, Hans

PA Roche Diagnostics G.m.b.H., Germany

SO Eur. Pat. Appl., 14 pp.

CODEN: EPXXDW

DT Patent

LA German

IC ICM C12M001-40

ICS C12P021-02; B01J019-24; B01J019-00  
 CC 9-1 (Biochemical Methods)  
 Section cross-reference(s): 3, 6, 7, 16

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1061128	A1	20001220	EP 1999-111762	19990618
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	CA 2340712	AA	20001228	CA 2000-2340712	20000610
	WO 2000078444	A2	20001228	WO 2000-EP5378	20000610
	WO 2000078444	A3	20010426		
	W: CA, JP, US				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	EP 1126912	A1	20010829	EP 2000-935201	20000610
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	JP 2003502146	T2	20030121	JP 2001-504498	20000610
PRAI	EP 1999-111762	A	19990618		
	WO 2000-EP5378	W	20000610		

AB The invention concerns a method and a device for high throughput biochem. **synthesis esp. cell free protein synthesis** that contains an outer vessel for the **transcription/translation** reagent supply; and an inner part that is a microtiter plate or similar multi-channel holder with semipermeable membrane bottom and used for the **transcription/translation** reaction components. The inner part contains 2-1000 wells; the vol. of the wells is 50 .mu.l - 10 mL. **Protein** -binding compds., e.g. avidin, streptavidin are immobilized on the inner walls of the wells for the facile purifn. of the **synthesized** tag-contg. **proteins**. The wells are capped or covered with foil. The system is on a shaker and/or stirred with magnetic stirrers inside the wells. The supply of reagents is maintained via diffusion. Thus two model **proteins, green fluorescence protein** and chloramphenicol-acetyltransferase were **synthesized** using **E.coli** lysate and 40 wells of a microtiterplate for each **protein**. **Transcription/translation** reagents, plasmid, **E.coli** lysate, T7 RNA polymerase were applied into the wells; the same reagent supply was used for both **proteins**. After 20 h of incubation at 30.degree.C and shaking, the **proteins** were measured in ech well.

ST high throughput **cell free protein synthesis** multichannel membrane device

IT **Proteins**, specific or class  
 RL: BPN (Biosynthetic preparation); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation)  
 (**green fluorescent**; method and multi-channel membrane device for high throughput **cell free protein synthesis**)

IT **Transcription, genetic**  
**Translation, genetic**  
 (in vitro; method and multi-channel membrane device for high throughput **cell free protein synthesis**)

IT **Escherichia coli**  
 (lysate; method and multi-channel membrane device for high throughput **cell free protein synthesis**)

IT Apparatus  
 Diffusion  
 Immobilization, biochemical  
 Microtiter plates  
 Plasmid **vectors**  
 Shaking apparatus

- (method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT **Fusion proteins (chimeric proteins)**  
RL: BPN (Biosynthetic preparation); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation)  
(method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT Avidins  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT Membranes, nonbiological  
(semipermeable; method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT **Proteins, general, preparation**  
RL: PUR (Purification or recovery); PREP (Preparation)  
(sepn.; method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT Mixers (processing apparatus)  
(stirrers; method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT 9040-07-7P, Chloramphenicol-acetyltransferase  
RL: BPN (Biosynthetic preparation); PRP (Properties); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation)  
(method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT 9013-20-1, Streptavidin  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)
- IT 9014-24-8  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(phage T7; method and multi-channel membrane device for high throughput  
**cell free protein synthesis**)

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Anon; PATENT ABSTRACTS OF JAPAN 1998, V1998(04)
- (2) Boehringer Ingelheim Kg; DE 3723004 A 1989
- (3) Boehringer Mannheim GmbH; EP 0894852 A 1999 HCAPLUS
- (4) July, B; US 5478730 A 1995 HCAPLUS
- (5) Kim, D; BIOTECHNOLOGY PROGRESS 1996, 5
- (6) Klaus Kuhn Konstruktion GmbH; DE 4237113 A 1994 HCAPLUS
- (7) Martin L Wolf & John Wilson; WO 9704074 A 1997
- (8) Rikagaku Kenkyusho & Sainikusu K K; JP 09322755 A 1997 HCAPLUS
- (9) Rikagaku Kenkyusho & Sainikusu K K; JP 09322755 A 1997 HCAPLUS
- (10) Rune, B; US 4642220 A 1987
- (11) Urban, S; US 5362624 A 1994 HCAPLUS

L60 ANSWER 29 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2000:895491 HCAPLUS

DN 135:222877

TI Structural genomics projects in Japan

AU Yokoyama, S.; Matsuo, Y.; Hirota, H.; Kigawa, T.; Shirouzu, M.; Kuroda, Y.; Kurumizaka, H.; Kawaguchi, S.; Ito, Y.; Shibata, T.; Kainosho, M.; Nishimura, Y.; Inoue, Y.; Kuramitsu, S.

CS Tsurumi-ku, 1-7-22 Suehiro-cho, RIKEN Genomic Sciences Center, Yokohama, 230-0045, Japan

SO Progress in Biophysics & Molecular Biology (2000), 73(5), 363-376  
CODEN: PBIMAC; ISSN: 0079-6107

PB Elsevier Science Ltd.  
DT Journal; General Review  
LA English  
CC 6-0 (General Biochemistry)  
AB A review with 22 refs. The "**Protein Folds Project**" and the "Structurome" project are described. Topics discussed include: **cell-free protein synthesis**; the large-scale NMR facility (NMR Farm); anal. of **protein function**; anal. of **protein-protein** interactions; selection of targets for structure detn.; identification of **protein domains**.  
ST review structural genomics project Japan; **protein** structure **function** detn review  
IT **Protein motifs**  
(identification; structural genomics projects in Japan)  
IT **Conformation**  
(**protein**, detn. by NMR; structural genomics projects in Japan)  
IT **Proteins**, general, biological studies  
RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
(structure-**function** detn.; structural genomics projects in Japan)  
RE.CNT 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE  
(1) Altschul, S; J Mol Biol 1990, V215, P403 HCAPLUS  
(2) Bartel, P; Nat genetic 1996, V12, P72 HCAPLUS  
(3) Brenner, S; Nucleic Acids Res 1999, V27, P251 HCAPLUS  
(4) Chothia, C; Nature 1992, V357, P543 MEDLINE  
(5) Davis, J; Prom Notes Mag 1996, V56, P14  
(6) Finley, R; Proc Natl Acad Sci USA 1994, V91, P12  
(7) Fromont-Racine, M; Nat genetic 1997, V16, P277 HCAPLUS  
(8) Gevaert, K; Electrophoresis 2000, V21, P1145 HCAPLUS  
(9) Hendrickson, W; EMBO J 1990, V9, P1665 HCAPLUS  
(10) Hwang, K; Nat Struct Biol 1999, V6, P691 HCAPLUS  
(11) Ito, T; Proc Natl Acad Sci USA 2000, V97, P1143 HCAPLUS  
(12) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS  
(13) Kigawa, T; J Biomol NMR 1995, V6, P129 HCAPLUS  
(14) Kigawa, T; The Abstract Book of the Seventh Annual Meeting of the Protein Engineering Society 1995, P48  
(15) Kim, D; Biotechnol Prog 1996, V12, P645 HCAPLUS  
(16) Kim, D; Eur J Biochem 1996, V239, P881 HCAPLUS  
(17) Kuramitsu, S; The Abstract Book of the seventh Annual Meeting of the Protein Engineering Society 1995, P48  
(18) Madin, K; Proc Natl Acad Sci USA 2000, V97, P559 HCAPLUS  
(19) McCraith, S; Proc Natl Acad Sci USA 2000, V97, P4879 HCAPLUS  
(20) Spirin, A; Science 1988, V242, P1162 HCAPLUS  
(21) Uetz, P; Nature 2000, V403, P623 HCAPLUS  
(22) Walhout, A; Science 2000, V287, P116 HCAPLUS  
  
L60 ANSWER 30 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2000:891657 HCAPLUS  
DN 134:4063  
TI Large-scale preparation of **proteins** by the **cell-free synthesis**  
AU **Kigawa, Takanori**  
CS RIKEN, Japan  
SO Seibutsu Butsuri (2000), 40(6), 391-394  
CODEN: SEBUAL; ISSN: 0582-4052  
PB Nippon Seibutsu Butsuri Gakkai  
DT Journal; General Review  
LA Japanese  
CC 16-0 (Fermentation and Bioindustrial Chemistry)

AB A review with 11 refs. on improvement of **protein** prepn. systems  
contg. **Escherichia coli** exts. and method for  
regioselective introduction of non-natural amino acids.

ST review **protein cell free** manuf; **Escherichia**  
ext **protein** manuf review

IT **Escherichia coli**  
Translation, genetic  
(large-scale prepn. of **proteins** by the **cell-**  
**free synthesis**)

IT **Proteins**, general, preparation  
RL: BMF (Bioindustrial manufacture); BIOL (Biological study); PREP  
(Preparation)  
(large-scale prepn. of **proteins** by the **cell-**  
**free synthesis**)

L60 ANSWER 31 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
AN 2000:510868 HCAPLUS  
DN 134:144124  
TI Large-scale purification of **recombinant green**  
**fluorescent protein** from **Escherichia**  
**coli**

AU Gonzalez, Daniel G.; Ward, William W.  
CS Department of Biochemistry and Microbiology, Cook College, Rutgers  
University, New Brunswick, NS, 08901-8525, Can.  
SO Methods in Enzymology (2000), 305 (Bioluminescence and Chemiluminescence,  
Pt. C), 212-223  
CODEN: MENZAU; ISSN: 0076-6879  
PB Academic Press  
DT Journal  
LA English  
CC 9-16 (Biochemical Methods)  
AB **Green fluorescent protein (GFP)**  
has been shown to form without the need for any cofactors or specific  
posttranslational processing from the native jellyfish in a **cell**  
**-free translation** system. It has been demonstrated,  
however, that mol. oxygen is necessary for chromophore formation. Its  
fluorescence properties allow for easy visualization and quantitation of  
**GFP** in a nondestructive manner. As a result, **GFP** has  
been used increasingly as a **fusion** partner for use in monitoring  
**expression**. The fluorescence from **GFP** may also be used  
to assay for a valuable **recombinant protein**  
nondestructively. **GFP** can be employed as a model  
**protein** to develop methods for **protein** purifn. It can  
also be utilized to demonstrate **protein** purifn. concepts in an  
instructional setting. Protocols used are hereby described. (c) 2000  
Academic Press.

ST purifn **recombinant green fluorescence**  
**protein Escherichia coli**

IT **Proteins**, specific or class  
RL: PUR (Purification or recovery); PREP (Preparation)  
(**green fluorescent**; large-scale purifn. of  
**recombinant green fluorescent**  
**protein** from **Escherichia coli**)

IT **Escherichia coli**  
(large-scale purifn. of **recombinant green**  
**fluorescent protein** from **Escherichia**  
**coli**)

IT **Proteins**, general, preparation  
RL: PUR (Purification or recovery); PREP (Preparation)  
(large-scale purifn. of **recombinant green**  
**fluorescent protein** from **Escherichia**  
**coli**)

RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD



RE

- (1) Anon; personal communication from D Davis
- (2) Chalfie, M; Science 1994, V263, P802 HCAPLUS
- (3) Cutler, M; Bioluminescence and Chemiluminescence: Fundamentals and Applied Aspects 1994, P383 HCAPLUS
- (4) Cutler, M; Ph D Thesis, Rutgers University 1995
- (5) Davis, D; Bioluminescence and Chemiluminescence: Fundamentals and Applied Aspects 1994, P596 HCAPLUS
- (6) Deschamps, J; Protein Express Purif 1995, V6, P555 HCAPLUS
- (7) Gonzalez, D; Green Fluorescent Protein: Properties, Applications, and Protocols 1998, P289
- (8) Inouye, S; FEBS Lett 1994, V341, P277 HCAPLUS
- (9) Johnson, B; Bio/Technology (London) 1994, V12, P1357 HCAPLUS
- (10) Kolb, V; Biotechnol Lett 1996, V18, P1447 HCAPLUS
- (11) Miller, J; Experiments in Molecular Genetics, 1972, P433
- (12) Morise, H; Biochemistry 1974, V13, P2656 HCAPLUS
- (13) Ormo, M; Science 1996, V273, P1392 HCAPLUS
- (14) Poppenborg, L; J Biotechnol 1977, V58, P79
- (15) Prasher, D; Gene 1992, V111, P229 HCAPLUS
- (16) Studier, F; J Mol Biol 1986, V189, P113 HCAPLUS
- (17) Studier, F; J Mol Biol 1991, V219, P37 HCAPLUS
- (18) Ward, W; Bioluminescence and Chemiluminescence: Basic Chemistry and Analytical Applications 1981, P235 HCAPLUS
- (19) Ward, W; Photochem Photobiol Rev 1979, V4, P1 HCAPLUS
- (20) Yang, F; Nature Biotechnol (London) 1996, V14, P1246 HCAPLUS

L60 ANSWER 32 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2000:336515 HCAPLUS.

DN 133:146435

TI Effects of anticodon 2'-O-methylations on tRNA codon recognition in an **Escherichia coli cell-free** translation

AU Satoh, Akira; Takai, Kazuyuki; Ouchi, Ryosuke; Yokoyama, Shigeyuki  
; Takaku, Hiroshi

CS Department of Industrial Chemistry, Chiba Institute of Technology, Chiba,  
275-0016, Japan

SO RNA (2000), 6(5), 680-686  
CODEN: RNARFU; ISSN: 1355-8382

PB Cambridge University Press

DT Journal

LA English

CC 6-2 (General Biochemistry)

AB The methylation of 2'-hydroxyl groups is one of the most common posttranscriptional modifications of naturally occurring stable RNA mols. Some tRNA species have a 2'-O-Me nucleoside at the first position of the anticodon, and it was suggested that this modification stabilizes the codon-anticodon duplex. However, no tRNA species have been found to have the modification at the second or third position of the anticodon. In the present study, we measured the effects of anticodon 2'-O-methylation on the codon-reading efficiencies of the anticodon variants of the unmodified forms of **Escherichia coli** tRNA<sup>Ser</sup>, using a **cell-free protein synthesis** assay.

The modification of C in the first position of the anticodon into 2'-O-methylcytidine increased the efficiency of reading the G-ending codon. On the other hand, the modifications of the second and/or third positions were detrimental to the codon-reading activity. Thus, 2'-hydroxyl groups at the second and third positions of the anticodon may have some role in the translation reaction, and this may be the reason why 2'-O-Me nucleosides are not found in these positions within natural tRNA species.

ST anticodon methylation tRNA codon recognition

IT Codons

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL

- (Biological study); PROC (Process)  
 (UCG and GAG recognition; **functional** role of anticodon  
 2'-O-methylation on tRNA codon recognition studied using  
 anticodon-modified derivs. of **synthetic** unmodified  
**Escherichia coli** tRNASer)
- IT Molecular recognition  
**Translation, genetic**  
**tRNA sequences**  
 (functional role of anticodon 2'-O-methylation on tRNA codon  
 recognition studied using anticodon-modified derivs. of  
**synthetic** unmodified **Escherichia coli**  
 tRNASer)
- IT Anticodons  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological  
 process); BSU (Biological study, unclassified); PRP (Properties); BIOL  
 (Biological study); PROC (Process)  
 (functional role of anticodon 2'-O-methylation on tRNA codon  
 recognition studied using anticodon-modified derivs. of  
**synthetic** unmodified **Escherichia coli**  
 tRNASer)
- IT Structure-activity relationship  
 (of tRNA anticodons; **functional** role of anticodon  
 2'-O-methylation on tRNA codon recognition studied using  
 anticodon-modified derivs. of **synthetic** unmodified  
**Escherichia coli** tRNASer)
- IT RNA  
 (processing; **functional** role of anticodon 2'-O-methylation on  
 tRNA codon recognition studied using anticodon-modified derivs. of  
**synthetic** unmodified **Escherichia coli**  
 tRNASer)
- IT tRNA  
 RL: BAC (Biological activity or effector, except adverse); BPN  
 (Biosynthetic preparation); BPR (Biological process); BSU (Biological  
 study, unclassified); BIOL (Biological study); PREP (Preparation); PROC  
 (Process)  
 (serine-specific UGA; **functional** role of anticodon  
 2'-O-methylation on tRNA codon recognition studied using  
 anticodon-modified derivs. of **synthetic** unmodified  
**Escherichia coli** tRNASer)
- IT 287121-69-1DP, derivs. contg. 2'-O-methylated and 2'-deoxy anticodon bases  
 287121-70-4DP, derivs. contg. 2'-O-methylated and 2'-deoxy anticodon bases  
 RL: BAC (Biological activity or effector, except adverse); BPN  
 (Biosynthetic preparation); BPR (Biological process); BSU (Biological  
 study, unclassified); BIOL (Biological study); PREP (Preparation); PROC  
 (Process)  
 (functional role of anticodon 2'-O-methylation on tRNA codon  
 recognition studied using anticodon-modified derivs. of  
**synthetic** unmodified **Escherichia coli**  
 tRNASer)
- IT 65-46-3, Cytidine 951-77-9, 2'-Deoxycytidine 2140-72-9,  
 2'-O-Methylcytidine  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
 (Biological study); PROC (Process)  
 (of anticodon; **functional** role of anticodon 2'-O-methylation  
 on tRNA codon recognition studied using anticodon-modified derivs. of  
**synthetic** unmodified **Escherichia coli**  
 tRNASer)

RE.CNT 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Ban, N; Nature 1999, V400, P841 HCAPLUS
- (2) Brimacombe, R; The many faces of RNA 1998, P41 HCAPLUS
- (3) Clemons, W; Nature 1999, V400, P833 HCAPLUS
- (4) Dao, V; Biochemistry 1992, V31, P11012 HCAPLUS

- (5) Dao, V; Proc Natl Acad Sci 1994, V91, P2125 HCAPLUS
- (6) Gabashvili, I; EMBO J 1999, V18, P6501 HCAPLUS
- (7) Glasser, A; FEBS Lett 1992, V314, P381 HCAPLUS
- (8) Horie, N; Biochemistry 1999, V38, P207 HCAPLUS
- (9) Inoue, H; Nucleic Acids Res 1987, V15, P6131 HCAPLUS
- (10) Kawai, G; Biochemistry 1992, V31, P1040 HCAPLUS
- (11) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (12) Lodmell, J; Science 1997, V277, P1262 HCAPLUS
- (13) Mendel, D; Annu Rev Biophys Biomol Struct 1995, V24, P435 HCAPLUS
- (14) Moore, P; Annu Rev Biophys Biomol Struct 1998, V27, P35 HCAPLUS
- (15) Murasugi, A; J Biochem 1978, V83, P1029 HCAPLUS
- (16) Noren, C; Science 1989, V244, P182 HCAPLUS
- (17) Ohtsuki, T; J Biochem 1996, V120, P1070 HCAPLUS
- (18) Pais de Barros, J; Nucleic Acids Res 1996, V24, P1489 HCAPLUS
- (19) Pillay, D; Nucleic Acids Res 1984, V12, P2997 HCAPLUS
- (20) Pixa, G; Biochem Biophys Res Commun 1983, V112, P578 HCAPLUS
- (21) Rothschild, K; Curr Opin Biotechnol 1999, V10, P64 HCAPLUS
- (22) Rozenski, J; Nucleic Acids Res 1999, V27, P196 HCAPLUS
- (23) Shimotono, K; J Mol Biol 1974, V86, P21 MEDLINE
- (24) Sprinzl, M; Nucleic Acids Res 1998, V26, P148 HCAPLUS
- (25) Takai, K; Biochem Biophys Res Commun 1999, V257, P662 HCAPLUS
- (26) Takai, K; FEBS Lett 1999, V447, P1 HCAPLUS
- (27) Takai, K; Nucleic Acids Res 1996, V24, P2894 HCAPLUS
- (28) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS
- (29) Yoshizawa, S; Science 1999, V285, P1722 HCAPLUS

L60 ANSWER 33 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2000:263884 HCAPLUS

DN 133:306799

TI Towards improved applications of **cell-free**

**protein** biosynthesis - the influence of mRNA structure and suppressor tRNAs on the efficiency of the system

AU Gerrits, Michael; Merk, Helmut; Stiege, Wolfgang; Erdmann, Volker A.

CS Institut fur Biochemie, Freie Universitat Berlin, Berlin, D-14195, Germany

SO NATO Science Series, 3: High Technology (1999), 70 (RNA Biochemistry and Biotechnology), 335-345

CODEN: NSSTFF; ISSN: 1388-6576

PB Kluwer Academic Publishers

DT Journal

LA English

CC 6-1 (General Biochemistry)

Section cross-reference(s): 7, 9

AB The **cell-free protein** biosynthesis has the potential to become a powerful technol. for the biochem. research in particular in the detn. of the structure and **function** of **proteins**. The no. of possible applications is rising with the obtainable yields and with the expanded feasibility of introducing modified amino acids into **proteins**. Here we describe the influence of two RNA **translation** components, the mRNA and the suppressor tRNA, on the efficiency of **protein** biosynthesis. It is shown that the rate limiting factor of the **cell-free translation** of the two **proteins** dihydrofolate reductase (DHFR) and fatty acid binding **protein** (FABP) is not the initiation or termination step. The efficiency of peptide bond formation in the nascent **protein** varies between the two genes but is independent on the size of the coding sequences. The poor **translation** of DHFR can be improved when its coding sequence is **fused** with a part of the more efficiently **translated** FABP gene. We compared different amber suppressor tRNAs on the level of **translational** efficiency and aminoacylation capacity. Our results show that in most cases the aminoacylation rate of the tRNAs is not the limiting factor of suppression. An *E. coli* tRNA<sup>Leu</sup>CUA exhibits the highest **translational** efficiency of the examd.

tRNAs. So this tRNA<sup>Leu</sup>CUA may be a starting point to construct more efficient tRNAs for the introduction of unnatural amino acids into **proteins** in the in vitro **translation** system by eliminating the **synthetase** mediated aminoacylation of the tRNA.

- ST **protein translation synthesis** mRNA structure  
tRNA suppressor; dihydrofolate reductase **translation synthesis** mRNA structure tRNA suppressor; FABP **protein translation synthesis** mRNA structure tRNA suppressor
- IT **Proteins**, specific or class  
RL: BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); SPN (Synthetic preparation); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); PROC (Process)  
(FABP (fatty acid-binding **protein**); towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)
- IT tRNA  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(amber suppressor; towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)
- IT tRNA  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(isoleucine-specific GAU; towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)
- IT tRNA  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(suppressor; towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)
- IT Aminoacylation  
**Translation, genetic**  
(towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)
- IT **Proteins**, general, biological studies  
RL: BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); SPN (Synthetic preparation); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); PROC (Process)  
(towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)
- IT mRNA  
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)
- IT 9002-03-3P, Dihydrofolate reductase  
RL: BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); SPN (Synthetic preparation); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); PROC (Process)  
(towards improved applications of **cell-free protein** biosynthesis - influence of mRNA structure and suppressor tRNAs on efficiency of system)

RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Cronenberger, J; J Mol Biol 1975, V95, P125 HCAPLUS
- (2) De Smit, M; Progr Nucl Acid Res Mol Biol 1990, V38, P1 HCAPLUS
- (3) Fuchs, U; FEBS Lett 1997, V414, P362 HCAPLUS
- (4) Gold, L; Annu Rev Microbiol 1981, V35, P365 HCAPLUS
- (5) Grosjean, H; J Mol Evolution 1978, V12, P113 HCAPLUS
- (6) Hall, M; Nature 1982, V295, P616 HCAPLUS
- (7) Haukanes, B; Biotechnology 1993, V11, P60 MEDLINE
- (8) Ikemura, T; J Mol Biol 1981, V151, P389 HCAPLUS
- (9) Karginov, V; Nucleic Acid Research 1997, V25, P3912 HCAPLUS
- (10) Kleina, L; J Mol Biol 1990, V213, P705 HCAPLUS
- (11) Laemmli, U; Nature 1970, V227, P680 HCAPLUS
- (12) Merk, H; to be published in J Biochem
- (13) Normanly, J; Nature 1986, V321, P213 HCAPLUS
- (14) Normanly, J; Proc Nat Acad Sci 1986, V83, P6548 HCAPLUS
- (15) Peterson, E; Biochemistry 1992, V31, P10380 HCAPLUS
- (16) Ringquist, S; Mol Microbiol 1992, V6, P1219 HCAPLUS
- (17) Robertson, S; J Am Chem Soc 1991, V113, P2722 HCAPLUS
- (18) Sambrook, J; Molecular cloning 1989
- (19) Sampson, J; Proc Nat Acad Sci 1988, V85, P1033 HCAPLUS
- (20) Stiege, W; J Biotechnol 1995, V41, P81 HCAPLUS
- (21) Triana-Alonso, F; J Biol Chem 1995, V270, P6298 HCAPLUS

L60 ANSWER 34 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 2000:183931 HCAPLUS

DN 132:319466

TI An easy **cell-free protein synthesis**  
system dependent on the addition of crude **Escherichia coli** tRNA

AU Kanda, Takayuki; Takai, Kazuyuki; Yokoyama, Shigeyuki; Takaku, Hiroshi

CS Department of Industrial Chemistry, Chiba Institute of Technology, Chiba, 275-0016, Japan

SO Journal of Biochemistry (Tokyo) (2000), 127(1), 37-41  
CODEN: JOBIAO; ISSN: 0021-924X

PB Japanese Biochemical Society

DT Journal

LA English

CC 9-16 (Biochemical Methods)

AB The **protein-synthesizing** S30 ext. of **Escherichia coli** contains tRNA, which limits its applications in **cell-free protein synthesis**. Here, we show that at least Arg- and Ser-acceptor activities can be removed from a std. S30 ext. by treatment with an immobilized RNase A resin. This RNase-treated ext. exhibits no **protein synthesis** activity, but regains it when supplied with crude **E. coli** tRNA and a small amt. of human placental RNase inhibitor. The **protein synthesis** is dependent on the addn. of tRNA in the presence of the RNase inhibitor. Chloramphenicol acetyltransferase was **synthesized** with this system and found to be active.

ST **protein synthesis** system **Escherichia** tRNAIT **Escherichia coli**

Translation, genetic

(easy **cell-free protein synthesis**system dependent on addn. of crude **Escherichia coli** tRNA)

IT tRNA

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(easy **cell-free protein synthesis**system dependent on addn. of crude **Escherichia coli**

tRNA)  
 IT 9040-07-7, Chloramphenicol acetyltransferase  
 RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL  
 (Biological study); FORM (Formation, nonpreparative)  
 (easy **cell-free protein synthesis**  
 system dependent on addn. of crude **Escherichia coli**  
 tRNA)

RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 RE

- (1) Brown, S; Cell 1987, V49, P825 HCAPLUS
- (2) Ehrenberg, M; Ribosomes and Protein Synthesis 1990, P101 HCAPLUS
- (3) Gesteland, R; Fed Proc 1965, V24, P293
- (4) Himeno, H; J Mol Biol 1997, V268, P803 HCAPLUS
- (5) Hohsaka, T; J Am Chem Soc 1996, V118, P9778 HCAPLUS
- (6) Hohsaka, T; J Am Chem Soc 1999, V121, P34 HCAPLUS
- (7) Kanamori, T; Proc Natl Acad Sci USA 1997, V94, P485 HCAPLUS
- (8) Kanamori, T; Proc Natl Acad Sci USA 1999, V96, P3634 HCAPLUS
- (9) Kanda, T; FEBS Lett 1998, V440, P273 HCAPLUS
- (10) Keiler, K; Science 1996, V271, P990 HCAPLUS
- (11) Kigawa, T; FEBS Lett 1999, V442, P15 HCAPLUS
- (12) Kigawa, T; J Biomol NMR 1995, V6, P129 HCAPLUS
- (13) Kim, D; Biotechnol Prog 1996, V12, P645 HCAPLUS
- (14) Kim, D; Eur J Biochem 1996, V239, P881 HCAPLUS
- (15) Kudlicki, W; J Mol Biol 1994, V244, P319 HCAPLUS
- (16) Ma, C; Biochemistry 1993, V32, P7939 HCAPLUS
- (17) Mendel, D; Annu Rev Biophys Biomol Struct 1995, V24, P435 HCAPLUS
- (18) Murakami, H; J Am Chem Soc 1998, V120, P7520 HCAPLUS
- (19) Muto, A; Trends Biochem Sci 1998, V23, P25 HCAPLUS
- (20) Nakamura, K; Eur J Biochem 1999, V259, P543 HCAPLUS
- (21) Nishimura, S; Biochim Biophys Acta 1967, V142, P133 HCAPLUS
- (22) Noren, C; Science 1989, V244, P182 HCAPLUS
- (23) Pavlov, M; Biochimie 1997, V79, P415 HCAPLUS
- (24) Pratt, J; Transcription and Translation 1984, P179 HCAPLUS
- (25) Shibata, T; J Biol Chem 1996, V271, P13162 HCAPLUS
- (26) Soberon, X; Gene 1980, V9, P287 HCAPLUS
- (27) Takai, K; Nucleic Acids Res 1996, V24, P2894 HCAPLUS
- (28) Wagner, E; Eur J Biochem 1982, V122, P193 HCAPLUS
- (29) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS
- (30) Zubay, G; J Mol Biol 1962, V4, P347 HCAPLUS

L60 ANSWER 35 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1999:156625 HCAPLUS

DN 130:219543

TI Large-scale **protein** preparation using the **cell-free synthesis**

AU **Kigawa, Takanori; Yabuki, Takashi; Yokoyama, Shigeyuki**

CS Cell. Signaling Lab., Inst. Phys. Chem. Res., Wako, 351-0198, Japan

SO Tanpakushitsu Kakusan Koso (1999), 44(4), 598-605

CODEN: TAKKAJ; ISSN: 0039-9450

PB Kyoritsu Shuppan

DT Journal; General Review

LA Japanese

CC 6-0 (General Biochemistry)

AB A review with 21 refs., on method for **cell-free synthesis** of **proteins** and its application to prepn. of amino acid-selective- and site-specific- labeled **proteins**, and development and application of large-scale **synthesis** system using dialysis. Method for tyrosine-selective labeling of Ras **protein** by **cell-free synthesis** is also discussed.

ST review **protein** prepn **cell free** system;  
 isotope labeling **protein synthesis** review

IT **Proteins**, specific or class

RL: BSU (Biological study, unclassified); RCT (Reactant); BIOL (Biological study); RACT (Reactant or reagent)  
(labeled; large-scale **protein** prepn. using **cell free synthesis**)

IT Genetic engineering

**Translation, genetic**

(large-scale **protein** prepn. using **cell free synthesis**)

IT **Proteins**, general, preparation

RL: BMF (Bioindustrial manufacture); BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

(large-scale **protein** prepn. using **cell free synthesis**)

L60 ANSWER 36 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1999:17197 HCAPLUS

DN 130:249005

TI **Cell-free** production and stable-isotope labeling of milligram quantities of **proteins**

AU **Kigawa, Takanori**; Yabuki, Takashi; Yoshida, Yasuhiko; Tsutsui, Michio; Ito, Yutaka; Shibata, Takehiko; **Yokoyama, Shigeyuki**

CS Cellular Signaling Laboratory, The Institute of Physical and Chemical Research (RIKEN), Wako, Saitama, 351-0198, Japan

SO FEBS Letters (1999), 442(1), 15-19

CODEN: FEBLAL; ISSN: 0014-5793

PB Elsevier Science B.V.

DT Journal

LA English

CC **9-8 (Biochemical Methods)**

AB We have improved the productivity of an **Escherichia coli** **cell-free protein synthesis** system.

First, creatine phosphate and creatine kinase were used as the energy source regeneration system, and the other components of the reaction mixt. were optimized. Second, the **E. coli** S30 **cell** ext. was condensed by dialysis against a polyethylene glycol soln. to increase the rate of **synthesis**. Third, during the **protein synthesis**, the reaction mixt. was dialyzed against a low-mol.-wt. substrate soln. to prolong the reaction. Thus, the yield of chloramphenicol acetyltransferase was raised to 6 mg/mL of reaction mixt. Stable-isotope labeling of a **protein** with <sup>13</sup>C/<sup>15</sup>N-labeled amino acids for NMR spectroscopy was achieved by this method.

ST **cell free** prodn isotope labeling **protein**

IT Dialysis

**Escherichia coli**

NMR (nuclear magnetic resonance)

(**cell-free** prodn. and stable-isotope labeling of milligram quantities of **proteins**)

IT **Proteins**, general, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(**cell-free** prodn. and stable-isotope labeling of milligram quantities of **proteins**)

IT Ras **proteins**

RL: BSU (Biological study, unclassified); BIOL (Biological study)

(**cell-free** prodn. and stable-isotope labeling of milligram quantities of **proteins**)

IT Isotopes

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(**cell-free** prodn. and stable-isotope labeling of milligram quantities of **proteins**)

IT 9040-07-7, Chloramphenicol acetyltransferase

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(**cell-free** prodn. and stable-isotope labeling of milligram quantities of **proteins**)

IT 67-07-2, Creatine phosphate 9001-15-4, Creatine kinase

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(**cell-free** prodn. and stable-isotope labeling of milligram quantities of **proteins**)

RE.CNT 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Choi, J; Anal Biochem 1996, V236, P82 HCAPLUS
- (2) Davis, J; Promega Notes Mag 1996, V56, P14
- (3) Ellman, J; Methods Enzymol 1991, V202, P301 HCAPLUS
- (4) Fujita-Yoshigaki, J; J Protein Chem 1992, V11, P731 HCAPLUS
- (5) Ha, J; Biochemistry 1989, V28, P8411 HCAPLUS
- (6) Ito, Y; Biochemistry 1997, V36, P9109 HCAPLUS
- (7) Kawarasaki, Y; Anal Biochem 1995, V226, P320 HCAPLUS
- (8) Kigawa, T; J Biomol NMR 1995, V6, P129 HCAPLUS
- (9) Kim, D; Biotechnol Prog 1996, V12, P645 HCAPLUS
- (10) Kim, D; Eur J Biochem 1996, V239, P881 HCAPLUS
- (11) Laemmli, U; Nature 1970, V227, P680 HCAPLUS
- (12) Muto, Y; J Biomol NMR 1993, V3, P165 HCAPLUS
- (13) Nakano, H; Biosci Biotechnol Biochem 1994, V58, P631 HCAPLUS
- (14) Nakano, H; J Biotechnol 1996, V46, P275 HCAPLUS
- (15) Odom, O; Anal Biochem 1997, V245, P249 HCAPLUS
- (16) Pelham, H; Eur J Biochem 1976, V67, P247 HCAPLUS
- (17) Pratt, J; No publication given 1984, P179 HCAPLUS
- (18) Ryabova, L; Anal Biochem 1995, V226, P184 HCAPLUS
- (19) Shaw, W; Methods Enzymol 1975, V43, P737 HCAPLUS
- (20) Sonar, S; Nature Struct Biol 1994, V1, P512 HCAPLUS
- (21) Spirin, A; Science 1988, V242, P1162 HCAPLUS
- (22) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS
- (23) Zawadzki, V; Nucleic Acids Res 1991, V19, P1948 HCAPLUS

L60 ANSWER 37 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1998:780895 HCAPLUS

DN 130:178868

TI Knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides

AU Kanda, Takayuki; Takai, Kazuyuki; Yokoyama, Shigeyuki; Takaku, Hiroshi

CS Faculty of Engineering, Department of Industrial Chemistry, Chiba Institute of Technology, Narashino, Chiba, 275-0016, Japan

SO FEBS Letters (1998), 440(3), 273-276

CODEN: FEBLAL; ISSN: 0014-5793

PB Elsevier Science B.V.

DT Journal

LA English

CC 6-1 (General Biochemistry)

AB Methods for the prepn. of an **Escherichia coli** tRNA mixt. lacking one or a few specific tRNA species can be the basis for future applications of **cell-free protein**

**synthesis**. We demonstrate here that virtually a single tRNA species in a crude **E. coli** tRNA mixt. can be knocked out by an antisense (complementary) oligodeoxyribonucleotide. One out of five oligomers complementary to tRNA<sup>Asp</sup> blocked the aspartylation almost completely, while minimally affecting the aminoacylation with other 13 amino acids tested. This 'knockout' tRNA behaved similarly to the untreated tRNA in a **cell-free** translation of an mRNA lacking Asp codons.

ST tRNA antisense oligodeoxyribonucleotide translation aminoacylation RNase H



- IT tRNA  
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(aspartic acid-specific; knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides)
- IT Aminoacylation  
**Translation, genetic**  
(knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides)
- IT DNA  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides)
- IT tRNA  
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides)
- IT Antisense oligonucleotides  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides)
- IT 9050-76-4, Ribonuclease H  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides)
- IT 220419-85-2 220419-98-7 220420-00-8 220420-01-9 220420-03-1  
RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
(knocking out a specific tRNA species within unfractionated **Escherichia coli** tRNA by using antisense (complementary) oligodeoxyribonucleotides)

RE.CNT 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD  
RE

- (1) Branch, A; Trends Biochem Sci 1998, V23, P45 HCAPLUS
- (2) Dong, H; J Mol Biol 1996, V260, P649 HCAPLUS
- (3) Ehrenberg, M; Ribosomes and Protein Synthesis 1990, P101 HCAPLUS
- (4) Hanes, J; Proc Natl Acad Sci USA 1997, V94, P4937 HCAPLUS
- (5) Harada, F; Biochem Biophys Res Commun 1972, V49, P1605 HCAPLUS
- (6) Hecht, S; J Biol Chem 1978, V253, P4517 HCAPLUS
- (7) Hohsaka, T; J Am Chem Soc 1996, V118, P9778 HCAPLUS
- (8) Hou, Y; Biochemistry 1996, V35, P15340 HCAPLUS
- (9) Komine, Y; J Mol Biol 1990, V212, P579 HCAPLUS
- (10) Ma, C; Biochemistry 1993, V32, P7939 HCAPLUS
- (11) Mendel, D; Annu Rev Biophys Biomol Struct 1995, V24, P435 HCAPLUS
- (12) Mirochnitchenko, O; Antisense Res Dev 1993, V3, P171 HCAPLUS
- (13) Murakami, H; J Am Chem Soc 1998, V120, P7520 HCAPLUS
- (14) Nishimura, S; Biochim Biophys Acta 1967, V142, P133 HCAPLUS
- (15) Noren, C; Science 1989, V244, P182 HCAPLUS
- (16) Takai, K; Nucleic Acids Res 1996, V24, P2894 HCAPLUS
- (17) Yabuki, T; J Biomol NMR 1998, V11, P295 HCAPLUS
- (18) Yokogawa, T; Nucleic Acids Res 1989, V17, P2623 HCAPLUS
- (19) Zubay, G; J Mol Biol 1962, V4, P347 HCAPLUS

L60 ANSWER 38 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1998:590136 HCAPLUS

DN 129:299303

TI A **cell-free protein synthesis**

system lacking specific tRNA species

AU Kanda, Takayuki; Takai, Kazuyuki; Yokoyama, Shigeyuki; Takaku, Hiroshi

CS Department of Industrial Chemistry, Chiba Institute of Technology, Tsudanuma, Narashino, Chiba, 275-0016, Japan

SO Nucleic Acids Symposium Series (1998), 39, 167-168

CODEN: NACSD8; ISSN: 0261-3166

PB Oxford University Press

DT Journal

LA English

CC 6-1 (General Biochemistry)

AB Prepn. of a **protein** with an unnatural or a labeled amino acid

residue was achieved by **cell-free protein**

**synthesis** supplied with an artificially-prepd. aminoacyl-tRNA that suppresses a stop codon. Sense codons cannot be used for this purpose, because a competing tRNA can also insert an unwanted natural amino acid at the position of the intended artificial amino acid. To utilize sense codons for the artificial amino acids, we have developed a strategy for the prepn. of **Escherichia coli** tRNA mixts. lacking the activity of a single tRNA species (tRNA<sup>Asp</sup> or tRNA<sup>Phe</sup>) based on oligonucleotide hybridization. In the present study, we demonstrated that these "knockout" tRNA mixts. cannot decode a codon corresponding to the inactivated tRNA, but can decode almost all of the other codons.

ST translation artificial amino acid tRNA

IT Codons

RL: BPR (Biological process); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); PROC (Process); USES (Uses)

(GAC and UUC; **cell-free protein**

**synthesis** system lacking specific tRNA species for utilizing sense codons for the artificial amino acids)

IT **Translation, genetic**

(**cell-free protein synthesis**

system lacking specific tRNA species for utilizing sense codons for the artificial amino acids)

IT tRNA

RL: BPR (Biological process); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); PROC (Process); USES (Uses)

(**cell-free protein synthesis**

system lacking specific tRNA species for utilizing sense codons for the artificial amino acids)

IT 56-87-1P, L-Lysine, preparation 60-18-4P, L-Tyrosine, preparation

72-19-5P, L-Threonine, preparation

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)

(**cell-free protein synthesis**

system lacking specific tRNA species for utilizing sense codons for the artificial amino acids)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Kanda, T; Nucleic Acids Res Sym Ser 1996, V35, P179

(2) Kanda, T; Nucleic Acids Res Sym Ser 1997, V37, P222

(3) Ma, C; Biochemistry 1993, V32, P7939 HCAPLUS

(4) Switzer, C; J Am Chem Soc 1989, V111, P8322 HCAPLUS

(5) Takai, K; Nucleic Acids Res 1996, V24, P2894 HCAPLUS

L60 ANSWER 39 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1998:477959 HCAPLUS  
 DN 129:200095  
 TI Dual amino acid-selective and site-directed stable-isotope labeling of the human c-Ha-Ras **protein** by **cell-free synthesis**  
 AU Yabuki, Takashi; **Kigawa, Takanori**; Dohmae, Naoshi; Takio, Koji; Terada, Tohru; Ito, Yutaka; Laue, Ernest D.; Cooper, Jonathan A.; Kainosho, Masatsune; **Yokoyama, Shigeyuki**  
 CS Cellular Signaling Laboratory, The Institute of Physical and Chemical Research (RIKEN), Saitama, 351-01, Japan  
 SO Journal of Biomolecular NMR (1998), 11(3), 295-306  
 CODEN: JBNME9; ISSN: 0925-2738  
 PB Kluwer Academic Publishers  
 DT Journal  
 LA English  
 CC 9-12 (Biochemical Methods)  
 Section cross-reference(s): 3, 7  
 AB We developed two methods for stable-isotope labeling of **proteins** by **cell-free synthesis**. Firstly, we applied **cell-free synthesis** to the dual amino acid-selective <sup>13</sup>C-<sup>15</sup>N labeling method, originally developed for in vivo systems by Kainosho and co-workers. For this purpose, we took one of the advantages of a **cell-free protein synthesis** system: the amino acid-selective stable-isotope labeling is **free** of the isotope scrambling problem. The targets of selective observation were Thr35 and Ser39 in the "effector region" (residues 32-40) of the Ras **protein** complexed with the Ras-binding domain of c-Raf-1 (Raf RBD) (the total mol. mass is about 30 kDa). Using a 15-mL **Escherichia coli cell-free** system, which was optimized to produce about 0.4 mg of Ras **protein** per 1-mL reaction, with 2 mg each of DL-[<sup>13</sup>C']proline and L-[<sup>15</sup>N]threonine, we obtained about 6 mg of Ras **protein**. As the Pro-Thr sequence is unique in the Ras **protein**, the Thr35 cross peak of the Ras.cntdot.Raf RBD complex was unambiguously identified by the 2D 1H-<sup>15</sup>N HNCQ expt. The Ser39 cross peak was similarly identified with the [<sup>13</sup>C']Asp/[<sup>15</sup>N]Ser-selectively labeled Ras **protein**. There were no isotope scrambling problems in this study. Secondly, we have established a method for producing a milligram quantity of site-specifically stable-isotope labeled **protein** by a **cell-free** system involving amber suppression. The **E. coli** amber suppressor tRNA<sup>Tyr</sup>CUA (25 mg) was prepd. by in vitro transcription with T7 RNA polymerase. We aminoacylated the tRNA<sup>Tyr</sup>CUA transcript with purified **E. coli** tyrosyl-tRNA **synthetase**, using 2 mg of L-[<sup>15</sup>N]tyrosine. In the gene encoding the Ras **protein**, the codon for Tyr32 was changed to an amber codon (TAG). This template DNA and the [<sup>15</sup>N]Tyr-tRNA<sup>Tyr</sup>CUA were reacted for 30 min in 30 mL of **E. coli cell-free** system. The subsequent purifn. yielded 2.2 mg of [<sup>15</sup>N]Tyr32-Ras **protein**. In the 1H-<sup>15</sup>N HSQC spectrum of the labeled Ras **protein**, only one cross peak was obsd., which was unambiguously assigned to Tyr32.  
 ST isotope labeling cHa Ras **protein**; **cell free synthesis** Ras **protein** Raf; HSQC NMR Ras labeled Raf RBD  
 IT Termination factors (**protein** formation)  
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (RF-1 (release factor 1); dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein** by **cell-free synthesis**)  
 IT Promoter (genetic element)  
 RL: BSU (Biological study, unclassified); BIOL (Biological study) (T7; dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein** by **cell-free synthesis**)

- IT Aminoacylation  
 Enzyme functional sites  
 Escherichia coli  
 Gel electrophoresis  
 Synthesis  
 Translation, genetic  
 (dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT Isotopes  
 RL: BSU (Biological study, unclassified); RCT (Reactant); BIOL (Biological study); RACT (Reactant or reagent)  
 (dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT Ras **proteins**  
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
 (dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT Transcription, genetic  
 (in vitro; dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT NMR spectroscopy  
 (multinuclear, HSQC spectrum; dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT Plasmids  
 (pK7-RasY32am and pK7-Ras; dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT Laser ionization mass spectrometry  
 (photodesorption, matrix-assisted; dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT Laser desorption mass spectrometry  
 (photoionization, matrix-assisted; dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT tRNA  
 RL: BPN (Biosynthetic preparation); PRP (Properties); RCT (Reactant); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)  
 (tyrosine-specific, **E.coli** amber suppressor tRNA<sup>tyr</sup>CUA; dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT 9014-24-8D, RNA polymerase, derivs.  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT 9023-45-4P, **Synthetase**, tyrosyl-transfer ribonucleate  
 RL: CAT (Catalyst use); PUR (Purification or recovery); PREP (Preparation); USES (Uses)  
 (dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free synthesis**)
- IT 59935-32-9, L-Serine-15N 80681-09-0, L-Threonine-15N 81201-97-0, L-Aspartic-1-13C acid 212258-93-0, Proline-carboxy-13C  
 RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)  
 (dual amino acid-selective and site-directed stable-isotope labeling of human c-Ha-Ras **protein by cell-free**)

**synthesis)**

IT 139691-76-2P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
(dual amino acid-selective and site-directed stable-isotope labeling of  
human c-Ha-Ras **protein** by **cell-free**

**synthesis)**

RE.CNT 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Akashi, S; J Mass Spectrom Soc Jpn 1996, V44, P269 HCAPLUS
- (2) Arata, Y; Methods Enzymol 1994, V239, P440 HCAPLUS
- (3) Bain, J; Biochemistry 1991, V30, P5411 HCAPLUS
- (4) Bodenhausen, G; Chem Phys Lett 1980, V69, P185 HCAPLUS
- (5) Delaglio, F; J Biomol NMR 1995, V6, P277 HCAPLUS
- (6) Ellman, J; J Am Chem Soc 1992, V114, P7959 HCAPLUS
- (7) Ellman, J; Methods Enzymol 1991, V202, P301 HCAPLUS
- (8) Grzesiek, S; J Am Chem Soc 1993, V115, P12593 HCAPLUS
- (9) Ha, J; Biochemistry 1989, V28, P8411 HCAPLUS
- (10) Ikura, M; Biochemistry 1990, V29, P4659 HCAPLUS
- (11) Ito, Y; Biochemistry 1997, V36, P9109 HCAPLUS
- (12) Johnson, B; J Biomol NMR 1994, V4, P603 HCAPLUS
- (13) Kainosho, M; Biochemistry 1982, V21, P6273 HCAPLUS
- (14) Kainosho, M; Biochemistry 1987, V26, P1068 HCAPLUS
- (15) Kainosho, M; J Mol Struct 1985, V126, P549 HCAPLUS
- (16) Kato, K; J Biochem (Tokyo) 1989, V105, P867 HCAPLUS
- (17) Ketchum, R; J Biomol NMR 1996, V8, P1 HCAPLUS
- (18) Kigawa, T; J Biomol NMR 1995, V6, P129 HCAPLUS
- (19) Kim, D; Eur J Biochem 1996, V239, P881 HCAPLUS
- (20) Kleina, L; J Mol Biol 1990, V213, P705 HCAPLUS
- (21) Liu, D; Proc Natl Acad Sci USA 1997, V94, P10092 HCAPLUS
- (22) Liu, X; Biophys Chem 1995, V56, P63 HCAPLUS
- (23) Lodder, M; J Org Chem 1997, V62, P778 HCAPLUS
- (24) McIntosh, L; Q Rev Biophys 1990, V23, P1 HCAPLUS
- (25) Milburn, M; Science 1990, V247, P939 HCAPLUS
- (26) Muto, Y; J Biomol NMR 1993, V3, P165 HCAPLUS
- (27) Nassar, N; Nat Struct Biol 1996, V3, P723 HCAPLUS
- (28) Nassar, N; Nature 1995, V375, P554 HCAPLUS
- (29) Noren, C; Science 1989, V244, P182 HCAPLUS
- (30) Normanly, J; J Mol Biol 1990, V213, P719 HCAPLUS
- (31) Pai, E; Nature 1989, V341, P209 HCAPLUS
- (32) Peersen, O; J Am Chem Soc 1992, V114, P4332 HCAPLUS
- (33) Pratt, J; Transcription and Translation 1984, P179 HCAPLUS
- (34) Robertson, S; J Am Chem Soc 1991, V113, P2722 HCAPLUS
- (35) Schagger, H; Anal Biochem 1987, V166, P368 MEDLINE
- (36) Schmitt, E; J Mol Biol 1993, V233, P615 HCAPLUS
- (37) Shulman, L; Prog Nucleic Acid Res Mol Biol 1991, V41, P23
- (38) Sonar, S; Nat Struct Biol 1994, V1, P512 HCAPLUS
- (39) Studier, F; Methods Enzymol 1990, V185, P60 HCAPLUS
- (40) Westler, W; J Am Chem Soc 1988, V110, P4093 HCAPLUS
- (41) Westler, W; J Am Chem Soc 1988, V110, P6256 HCAPLUS
- (42) Yamasaki, K; J Biomol NMR 1992, V2, P71 HCAPLUS
- (43) Zawadzki, V; Nucleic Acids Res 1991, V19, P1948 HCAPLUS

L60 ANSWER 40 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1998:295237 HCAPLUS

DN 129:50985

TI Removing tRNA from a **cell-free protein****synthesis** system for use in **protein** productionAU Kanada, Takayuki; Takai, Kazuyuki; Yokoyama, Shigeyuki; Takaku,  
HiroshiCS Dep. of Industrial Chemistry, Chiba Institute of Technology, Narashino,  
275, JapanSO Nucleic Acids Symposium Series (1997), 37(Symposium on Nucleic Acids  
Chemistry, 1997), 319-320

CODEN: NACSD8; ISSN: 0261-3166

PB Oxford University Press

DT Journal

LA English

CC 6-2 (General Biochemistry)

AB The **cell-free** system for biosynthesis of**proteins** is becoming an important tool for **protein**engineering. In particular, introduction of the unnatural amino acids is achieved though **cell-free protein****synthesis** with the use of chem. acylated tRNA that recognizes a specific codon. In the original method, however, it was difficult to control the system through changing tRNA compn., as the endogenous tRNAs are involved in the reaction. Thus, in the present study, we**digested** the tRNA within **Escherichia coli** S30ext. with resin-bound RNase A, and estd. the **protein****synthesis** activity. It was revealed that this **digestion**process does not damage the activity, if a protease inhibitor, phenylmethylsulfonyl fluoride (PMSF), is present in the **digestion** reaction.ST tRNA translation **protein** engineering RNase AIT **Protein** engineering**Translation, genetic**(removing tRNA from a **cell-free protein****synthesis** system for use in **protein** prodn.)

IT tRNA

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(removing tRNA from a **cell-free protein****synthesis** system for use in **protein** prodn.)

IT 9001-99-4, Ribonuclease

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(A; removing tRNA from a **cell-free protein****synthesis** system for use in **protein** prodn.)

IT 329-98-6, Phenylmethylsulfonyl fluoride

RL: BOC (Biological occurrence); BSU (Biological study, unclassified);

BIOL (Biological study); OCCU (Occurrence)

(removing tRNA from a **cell-free protein****synthesis** system for use in **protein** prodn.)

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

(1) Kanda, T; Nucleic Acids Res Sym Ser 1996, V35, P179

(2) Ma, C; Biochemistry 1993, V32, P7939 HCAPLUS

(3) Noren, C; Science 1989, V244, P182 HCAPLUS

(4) Pratt, J; Transcription and translation 1984, P179 HCAPLUS

(5) Takai, K; Nucleic Acids Res 1996, V24, P2894 HCAPLUS

L60 ANSWER 41 OF 50 .HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1997:356920 HCAPLUS

DN 127:106257

TI Correspondence: **Cell-free expression** of a**GFP fusion protein** allows quantitation in

vitro and in vivo

AU Kahn, Theodore W.; Beachy, Roger N.; Falk, Matthias M.

CS Department Cell Biology, Scripps Research Institute, La Jolla, CA, 92037, USA

SO Current Biology (1997), 7(4), R207-R208

CODEN: CUBLE2; ISSN: 0960-9822

PB Current Biology

DT Journal

LA English

CC 9-16 (Biochemical Methods)

AB When a **GFP (green fluorescent**

**protein) fusion protein** is translated in vitro under std. conditions, the **GFP** portion **folds** efficiently and becomes fluorescent. This provides a convenient method for monitoring in vitro translation efficiency of a **fusion protein**, and to screen for improving mutants of **GFP**.

ST cell **GFP fusion protein**

IT **Proteins**, specific or class

RL: BPR (Biological process); BSU (Biological study, unclassified); BUU (Biological use, unclassified); BIOL (Biological study); PROC (Process); USES (Uses)

(**green fluorescent, fusion protein**; correspondence: **cell-free expression of a GFP fusion protein** allows quantitation in vitro and in vivo)

L60 ANSWER 42 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1997:228751 HCAPLUS

DN 126:313776

TI **Protein** engineering of de novo **protein** with predesigned structure and activity

AU Dolgikh, Dmitry A.; Gabrielian, Andrei E.; Uversky, Vladimir N.; Kirpichnikov, Michael P.

CS Institute of Molecular Biology, Russian Academy of Sciences, Moscow, 117984, Russia

SO Applied Biochemistry and Biotechnology (1997), Volume Date 1996, 61(1/2, Biocatalysis-95), 85-96

CODEN: ABIBDL; ISSN: 0273-2289

PB Humana

DT Journal

LA English

CC 6-3 (General Biochemistry)

Section cross-reference(s): 3

AB The de novo **protein** albebetin has been engineered (J. Mol. Biol. 1992, 225, 927-931) to form a predesigned tertiary **fold** that has not yet been obsd. in natural **proteins**. Anal. of albebetin **expressed** in a **cell-free** system and in

**Escherichia coli** revealed its compactness, relative stability, and the secondary structure close to the predesigned one. The blast-transforming biol. activity of human interferon was grafted to albebetin by attachment of an eight amino acid interferon fragment to the N-terminus of albebetin next to its first methionine residue. The chimeric **protein** was **expressed** in a wheat germ **cell-free translation** system and tested for its structural properties, receptor binding, and biol. activity. According to the tests, albebetin incorporating the active interferon fragment has a compact and relatively stable structure, and binds the murine thymocyte receptor effectively. It activates the blast transformation reaction of thymocyte **cells** even more efficiently than human interferon at low concns.

ST albebetin interferon conjugate **recombinant** receptor binding

IT **Proteins**, specific or class

RL: BAC (Biological activity or effector, except adverse); BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)

(albebetin, **fusion protein** with interferon .alpha.2 fragment; **protein** engineering of de novo **protein** with predesigned structure and activity)

IT **Escherichia coli**

(albebetin-interferon conjugate prodn. with; **protein** engineering of de novo **protein** with predesigned structure and activity)

IT **DNA sequences**

(for interferon .alpha.2-albebetin **fusion protein**)

IT **Protein sequences**  
(of interferon .alpha.2-albebetin **fusion protein**)

IT Interferons  
RL: BAC (Biological activity or effector, except adverse); BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)  
(.alpha.2, **fusion protein** with albebetin;  
**protein** engineering of de novo **protein** with  
predesigned structure and activity)

IT 148522-27-4P  
RL: BAC (Biological activity or effector, except adverse); BPN (Biosynthetic preparation); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PREP (Preparation)  
(amino acid sequence; **protein** engineering of de novo  
**protein** with predesigned structure and activity)

IT 189355-70-2  
RL: BUU (Biological use, unclassified); PRP (Properties); BIOL (Biological study); USES (Uses)  
(nucleotide sequence; **protein** engineering of de novo  
**protein** with predesigned structure and activity)

L60 ANSWER 43 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1996:554280 HCAPLUS

DN 125:188549

TI Codon-reading specificity of an unmodified form of **Escherichia coli** tRNA<sup>1</sup>Ser in **cell-free protein synthesis**

AU Takai, Kazuyuki; Takaku, Hiroshi; Yokoyama, Shigeyuki

CS Dep. Biophys. Biochem., Univ. Tokyo, Tokyo, 113, Japan

SO Nucleic Acids Research (1996), 24(15), 2894-2899

CODEN: NARHAD; ISSN: 0305-1048

PB Oxford University Press

DT Journal

LA English

CC 6-1 (General Biochemistry)

Section cross-reference(s): 3

AB Unmodified tRNA mols. are useful for many purposes in **cell-free protein** biosynthesis, but there is little information about how the lack of tRNA posttranscriptional modifications affects the coding specificity for synonymous codons. In the present study, we prepd. an unmodified form of **Escherichia coli** tRNA<sup>1</sup>Ser, which originally has the cmo5UGA anticodon (cmo5U = uridine 5-oxyacetic acid) and recognizes the UCU, UCA and UCG codons. The codon specificity of the unmodified tRNA was tested in a **cell-free protein synthesis** directed by designed mRNAs under competition conditions with the parent tRNA<sup>1</sup>Ser. It was found that the unmodified tRNA with the UGA anticodon recognizes the UCA codon nearly as efficiently as the modified tRNA. The unmodified tRNA recognized the UCU codon with low, but detectable efficiency, whereas no recognition of the UCC and UCG codons was detected. Therefore, the absence of modifications makes this tRNA more specific to the UCA codon by remarkably reducing the efficiencies of wobble reading of other synonymous codons, without a significant decrease in the UCA reading efficiency.

ST codon specificity **Escherichia** tRNA<sup>1</sup>Ser **protein synthesis**

IT **Escherichia coli**  
**Translation, genetic**  
(codon-reading specificity of an unmodified form of **Escherichia coli** tRNA<sup>1</sup>Ser in **cell-free protein synthesis**)

IT Codon  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(codon-reading specificity of an unmodified form of **Escherichia**



**coli tRNA<sup>1</sup>Ser in cell-free protein synthesis)**

- IT Ribonucleic acids, transfer  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (serine-specific 1, codon-reading specificity of an unmodified form of **Escherichia coli tRNA<sup>1</sup>Ser in cell-free protein synthesis)**)
- L60 ANSWER 44 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1996:512916 HCAPLUS  
 DN 125:162430  
 TI A highly efficient **cell-free protein synthesis** system from **Escherichia coli**  
 AU Kim, Dong-Myung; Kigawa, Takanori; Choi, Cha-Yong; Yokoyama, Shigeyuki  
 CS Interdisciplinary Program Biochemical Eng. and Technology, Seoul Natl. Univ., S. Korea  
 SO European Journal of Biochemistry (1996), 239(3), 881-886  
 CODEN: EJBCAI; ISSN: 0014-2956  
 PB Springer  
 DT Journal  
 LA English  
 CC 9-2 (Biochemical Methods)  
 AB The authors modified a **cell-free** coupled transcription/translation system from **Escherichia coli** with the T7 phage RNA polymerase, and achieved a productivity as high as 0.4 mg **protein**/mL reaction mixt. First, the authors found that the optimal concns. of phosphoenolpyruvate and poly(ethylene glycol) are interdependent; higher concns. of the former should be used at higher concns. of the latter. Second, the use of a condensed 30,000 g **cell ext.**, in place of the conventional one, significantly increased the initial rate of **protein synthesis**. This phenomenon was demonstrated to be due to a reason other than elimination of inhibitory mol(s). from the ext. For this system with the condensed ext., the phosphoenolpyruvate and poly(ethylene glycol) concns. were again co-optimized, resulting in prodn. of chloramphenicol acetyltransferase at a productivity of 0.3 mg/mL. Finally, the productivity was further increased up to 0.4 mg/mL, by supplementation of the pool of amino acids. This improved **cell-free protein synthesis** system is superior in productivity to any other **cell-free** systems reported so far, including the continuous-flow **cell-free** system.
- ST **cell free** translation system **Escherichia**  
 IT **Escherichia coli**  
     **Translation, genetic**  
     (highly efficient **cell-free protein synthesis** system from **Escherichia coli**)
- IT Virus, bacterial  
     (T7, highly efficient **cell-free protein synthesis** system from **Escherichia coli**)
- IT 9040-07-7P, Chloramphenicol acetyltransferase  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
     (highly efficient **cell-free protein synthesis** system from **Escherichia coli**)
- IT 138-08-9, Phosphoenolpyruvic acid 9014-24-8, RNA polymerase 25322-68-3  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
     (highly efficient **cell-free protein synthesis** system from **Escherichia coli**)

AN 1995:871566 HCAPLUS  
 DN 123:250460  
 TI **Cell-free synthesis** and amino acid-selective  
 stable isotope labeling of **proteins** for NMR analysis  
 AU **Kigawa, Takanori**; Muto, Yutaka; **Yokoyama, Shigeyuki**  
 CS Dep. Biophys. Biochem. Sch. Sci., Univ. Tokyo, Tokyo, 113, Japan  
 SO Journal of Biomolecular NMR (1995), 6(2), 129-34  
 CODEN: JBNME9; ISSN: 0925-2738  
 PB ESCOM  
 DT Journal  
 LA English  
 CC **9-5 (Biochemical Methods)**  
 Section cross-reference(s): 6, 77  
 AB For the application of multidimensional NMR spectroscopy to larger  
**proteins**, it would be useful to perform selective labeling of one  
 of the 20 amino acids. For some amino acids, however, amino acid metab.  
 drastically reduces the efficiency and selectivity of labeling in in vivo  
**expression** systems. In the present study, a **cell-**  
**free protein synthesis** system was optimized,  
 so that highly efficient and selective stable isotope labeling of  
**proteins** can be achieved in the absence of amino acid metab. The  
 productivity of the **E. coli cell-**  
**free** coupled transcription-translation system was first improved,  
 by about 5-fold, by using the T7 RNA polymerase for  
 transcription and also by improving the translation conditions. Thus,  
 about 0.1 mg **protein** per 1 mL reaction mixt. was  
**synthesized**. Then, this improved **cell-free**  
 system was used for Asp- or Ser-selective <sup>15</sup>N-labeling of the human  
 c-Ha-Ras **protein**. With a 15 mL **cell-free**  
 reaction, using <1 mg of <sup>15</sup>N-labeled amino acid, 1 mg of the Ras  
**protein** was obtained. <sup>1</sup>H-<sup>15</sup>N HSQC expts. confirmed that the Ras  
**protein** was efficiently labeled with high selectivity. These  
 results indicate that this **cell-free protein**  
**synthesis** system is useful for NMR studies.  
 ST **protein** stable isotope labeling NMR analysis; **cell**  
**free protein synthesis** isotope labeling;  
 nitrogen 15 labeling cHaRas **protein**  
 IT **Escherichia coli**  
 Nuclear magnetic resonance spectrometry  
 (**cell-free synthesis** and amino  
 acid-selective stable isotope labeling of **proteins** for NMR  
 anal.)  
 IT **Proteins**, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (**cell-free synthesis** and amino  
 acid-selective stable isotope labeling of **proteins** for NMR  
 anal.)  
 IT G **proteins** (guanine nucleotide-binding **proteins**)  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (p21c-Ha-ras, **cell-free synthesis** and  
 amino acid-selective stable isotope labeling of **proteins** for  
 NMR anal.)  
 IT Isotope indicators  
 (stable, **cell-free synthesis** and amino  
 acid-selective stable isotope labeling of **proteins** for NMR  
 anal.)  
 IT 14390-96-6, Nitrogen 15, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (**cell-free synthesis** and amino  
 acid-selective stable isotope labeling of **proteins** for NMR  
 anal.)

AN 1994:4060 HCAPLUS  
 DN 120:4060  
 TI Labeling of **proteins** in **cell-free**  
**protein synthesis** system containing non-natural amino  
 acid substrates  
 IN **Yokoyama, Shigeyuki; Kikawa, Takanori; Tejima,**  
 Munehiro  
 PA Nippon Oxygen Co Ltd, Japan  
 SO Jpn. Kokai Tokkyo Koho, 9 pp.  
 CODEN: JKXXAF

DT **Patent**  
 LA Japanese  
 IC ICM C12P021-00  
 ICS C12P021-00  
 ICA C12N015-09  
 CC **9-14 (Biochemical Methods)**  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 05236987	A2	19930917	JP 1991-208321	19910820
	JP 3317983	B2	20020826		
PRAI	JP 1991-208321		19910820		

AB A method for labeling **proteins** with non-natural amino acid  
 substrates in a **cell-free protein**  
**synthesis** system and a kit for the method are provided. The  
 system employs codon-specific isoaccepting tRNA and the corresponding  
 non-natural amino acids so that the labeling can be directed to specific  
 triplet codon(s). The non-natural amino acids can be the natural amino  
 acids having atoms substituted with a stable isotope or a radioisotope,  
 enantiomeric side chains, or substituted side chains that cause changes of  
 properties (e.g. mol. size, reactivity, hydrophobicity, fluorescence,  
 hydrogen bonding, etc). A kit for **cell-free**  
**protein synthesis** contg. ribosomes, tRNAs,  
**enzymes**, isoaccepting tRNA bound with (non-)natural amino acids,  
 and ribonucleotides and the **proteins** prep'd. with the method are  
 claimed. Prepn. of chlroamphenicol acetyl transferase (CAT) using  
**Escherichia coli** S30 ext. was shown, in which natural  
 arginyl tRNA1, tRNA2, and tRNA3 and 15N-labeled tRNA4 were used. The  
 resultant CAT contained 15N-Arg at position 213, but not the rest of Arg  
 (at positions 18, 65, 74, and 124).

ST **protein** labeling artificial amino acid; tRNA isoaccepting  
**protein** labeling

IT Isotopes  
 Radioelements, uses  
 RL: ANST (Analytical study)  
 (amino acids labeled with, for **protein** labeling, isoaccepting  
 tRNA in **cell-free protein**  
**synthesis** system in relation to)

IT **Translation, genetic**  
 (**cell-free**, isoaccepting tRNA in, for  
**protein** labeling)

IT Ribonucleic acids, transfer  
 RL: ANST (Analytical study)  
 (isoaccepting, in codon-specific labeling of **proteins**)

IT **Proteins**, biological studies  
 RL: BIOL (Biological study)  
 (labeling of, codon-specific, isoaccepting tRNA in)

IT Amino acids, uses  
 RL: USES (Uses)  
 (**proteins** labeling with, isoaccepting tRNA in)

IT Isomerism and Isomers  
 (optical, amino acids labeled with, for **protein** labeling,  
 isoaccepting tRNA in **cell-free protein**

**synthesis** system in relation to)

L60 ANSWER 47 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1994:4034 HCAPLUS  
 DN 120:4034  
 TI Labeling of **proteins** with stable radioisotopes and kits for labeling  
 IN **Yokoyama, Shigeyuki; Kikawa, Takanori; Tejima, Munehiro**  
 PA Nippon Oxygen Co Ltd, Japan  
 SO Jpn. Kokai Tokkyo Koho, 6 pp.  
 CODEN: JKXXAF  
 DT **Patent**  
 LA Japanese  
 IC ICM C12P021-00  
 ICA C12N015-09  
 CC **9-8 (Biochemical Methods)**  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 05236986	A2	19930917	JP 1991-208322	19910820
	JP 3145431	B2	20010312		
PRAI	JP 1991-208322		19910820		

AB A method for labeling **proteins** with stable radioisotopes in a **cell-free, protein synthetic** system is described. The system is continuously supplied with amino acid substrates labeled with radioisotopes such as <sup>13</sup>C and <sup>15</sup>N and the product can be almost 100% labeled. It also contains the amino acid **synthesis** system, an inhibitor for amino acid metab., and ribonucleotides. Other **enzymes** from the **cell** origins are removed by, e.g., monoclonal antibodies. The method was demonstrated by **synthesis** of <sup>15</sup>N-labeled CAT using the **Escherichia coli** S-30 ext.

ST **cell free protein synthesis**  
 radioisotope labeling

IT **Proteins**, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (labeling of, with radioisotopes, **cell-free protein synthetic** system for)

IT Radioelements, reactions  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (**protein** labeling with, **cell-free protein synthetic** system for)

L60 ANSWER 48 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN  
 AN 1992:610746 HCAPLUS  
 DN 117:210746  
 TI **Cell-free**, continuous **synthesis** of polypeptide  
 IN **Yokoyama, Shigeyuki; Endo, Yaeta; Kikawa, Takanori**  
 PA Japan  
 SO Jpn. Kokai Tokkyo Koho, 7 pp.  
 CODEN: JKXXAF  
 DT **Patent**  
 LA Japanese  
 IC ICM C12P021-00  
 CC 16-4 (Fermentation and Bioindustrial Chemistry)  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 04200390	A2	19920721	JP 1990-334103	19901130
PRAI	JP 1990-334103		19901130		

AB A **cell-free**, ribosome-contg., continuous system for

**synthesis** of polypeptide (excluding chem. **synthesis**) from mRNA is disclosed. In this system, the reaction substrates, e.g., ATP, GTP, amino acids, etc., are continuously supplied to the reaction chamber having minimal air content and the reaction mixt. is pumped through an ultrafiltration app. to sep. polypeptide products from the low-mol. substrates. The process minimizes foaming and therefore **protein** denaturation as compared to that of prior art.

**Synthesis** of chloramphenicol acetyltransferase (CAT) was performed continuously for 17 h with a good yield.

ST **cell free** continuous polypeptide **synthesis**;  
foaming **free protein synthesis cell**  
**free**

IT Foaming  
(**cell-free** and continuous **protein**  
**synthesis** system low in)

IT **Translation, genetic**  
(**cell-free** and continuous system for, low-foaming)

IT Air  
(min., in **cell-free** and continuous system for  
**protein synthesis, protein** denaturation  
minimization in relation to)

IT Denaturation  
(**protein, foaming** in, **cell-free** and  
continuous system for **protein synthesis** avoiding,  
min. air content in relation to)

IT Peptides, preparation  
**Proteins, preparation**  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(**synthesis** of, **cell-free** and continuous  
system low in foaming for)

IT 9040-07-7  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(**synthesis** of, **cell-free** and continuous  
system low in foaming for)

L60 ANSWER 49 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN

AN 1991:581406 HCAPLUS

DN 115:181406

TI A continuous **cell-free protein**  
**synthesis** system for coupled transcription-translation

AU Kigawa, Takanori; Yokoyama, Shigeyuki

CS Fac. Sci., Univ. Tokyo, Tokyo, 113, Japan

SO Journal of Biochemistry (Tokyo, Japan) (1991), 110(2), 166-8  
CODEN: JOBIAO; ISSN: 0021-924X

DT Journal

LA English

CC 16-1 (Fermentation and Bioindustrial Chemistry)

Section cross-reference(s): 9

AB A continuous **cell-free protein**

**synthesis** system (Spirin, A. S. et al., 1988) was modified so as to be suitable for coupled transcription-translation, a process useful for obtaining products of **cloned** genes or cDNAs. A reaction chamber equipped with an ultrafiltration membrane was newly designed and an HPLC pump was used to supply a low-mol.-wt. substrate soln. at a const. rate to the viscous reaction mixt. in the chamber. By using an **Escherichia coli** S30 ext. in this modified flow system (1 mL), coupled transcription-translation could be continuously performed for 17 h, the **synthesized** chloramphenicol acetyltransferase (.simeq.0.1 mg) being subsequently eluted through the chamber membrane and then purified.

ST reactor continuous **protein synthesis** system; coupled  
transcription translation **Escherichia; cell free**  
system transcription translation

- IT **Escherichia coli**  
(continuous **cell-free protein synthesis** system for coupled translation and transcription based on exts. on)
- IT **Protein formation**  
(continuous **cell-free** system for, for coupled transcription-translation)
- IT Deoxyribonucleic acids  
RL: BIOL (Biological study)  
(transcription of, translation coupled with, continuous **cell-free protein synthesis** system for)
- IT Ribonucleic acids, messenger  
RL: BIOL (Biological study)  
(translation of, transcription coupled with, continuous **cell-free protein synthesis** system for)
- IT Reactors  
(ultrafiltration membrane combined with, in continuous **cell-free protein synthesis** system for coupled transcription-translation)
- IT Filtering materials  
(ultra-, membranes, in reaction chamber, for continuous **cell-free protein synthesis** system for coupled transcription-translation)
- IT 9040-07-7P, Chloramphenicol acetyltransferase  
RL: PREP (Preparation)  
(prepn. of, in continuous **cell-free protein synthesis** system)
- L60 ANSWER 50 OF 50 HCAPLUS COPYRIGHT 2003 ACS on STN
- AN 1991:178686 HCAPLUS
- DN 114:178686
- TI Insulin and IGF-I induce pronounced hypertrophy of skeletal myofibers in tissue culture
- AU Vandeburgh, Herman H.; Karlisch, Patricia; Shansky, Janet; Feldstein, Rebecca
- CS Dep. Pathol., Brown Univ., Providence, RI, 02906, USA
- SO American Journal of Physiology (1991), 260(3, Pt. 1), C475-C484  
CODEN: AJPHAP; ISSN: 0002-9513
- DT Journal
- LA English
- CC 2-6 (Mammalian Hormones)
- AB Skeletal myofibers differentiated from primary avian myoblasts in tissue culture can be maintained in pos. N balance in a defined serum-**free** medium for at least 6-7 days when embedded in a 3-**dimensional** collagen gel matrix. Incubation of established myofiber cultures for 3-7 days with insulin (1 .mu.M) or insulin-like growth factor I (IGF-I, 32 nM) stimulates both **cell** hyperplasia and myofiber hypertrophy. Mean myofiber diam. increases 71-98%. Insulin-like growth factor II stimulates **cell** hyperplasia but not myofiber hypertrophy. **Cell** growth results from a 42-62% increase in total **protein synthesis** and a 28-38% decrease in **protein** degrdn. Myosin heavy-chain content increases 183-258% because of a 55% stimulation of myosin **synthesis** and 33-61% inhibition of degrdn. Assocd. with myofiber hypertrophy is a 87-148% increase in the no. of myofiber nuclei per unit myofiber length. The results indicate that insulin and IGF-I, but not IGF-II, can induce rapid myofiber hypertrophy in vitro, most likely by stimulating myoblast proliferation and/or **fusion** to established myofibers.
- ST insulin skeletal myofiber hypertrophy; insulin growth factor myofiber hypertrophy
- IT Myosins  
RL: BIOL (Biological study)

(heavy chain, formation of, by muscle myofibers in culture, insulin and insulin-like growth factor I stimulation of)

IT **Proteins**, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(metab. of, by muscle myofibers in culture, insulin and insulin-like growth factor I inhibition of)

IT **Protein formation**

(muscle cell growth stimulation by insulin and insulin-like growth factor I mediation by, in culture)

IT **Muscle**

(fiber, hypertrophy of, in culture, insulin and insulin-like growth factor I induction of)

IT 67763-97-7, Insulin-like growth factor II

RL: PROC (Process)

(muscle myofiber cell hyperplasia stimulation of, in culture)

IT 9004-10-8, Insulin, biological studies 67763-96-6, Insulin-like growth factor I

RL: BIOL (Biological study)

(muscle myofiber hypertrophy induction by, in culture)